

MICHIGAN INFANT DEATH STATISTICS
January 1 through December 31, 2015

LANSING, MICHIGAN
MARCH 2017



Rick Snyder, Governor
Nick Lyon, Director

Acknowledgments

DIVISION FOR VITAL RECORDS AND HEALTH STATISTICS

Glenn Copeland, M.B.A. State Registrar

VITAL RECORDS & HEALTH DATA DEVELOPMENT SECTION

Tami Weaver, B.A., Manager

AUTHORS

Glenn Radford, M.A., Statistician
Lindsey Myers, B.A., Data Analyst

The Division for Vital Records and Health Statistics gratefully acknowledges the contributions of: Kay Bertrau, B.A., Thu Le, Ph.D., Kyle Johnson and Kim Essex.



Table Of Contents

Summary	1
---------------	---

Selected Infant Death Statistics, Numbers, Averages & Rates

Table 1 Number of Infant Deaths, Live Births & Infant Death Rates for Michigan Residents, 1970-2015	4
Figure 1 Infant Mortality Rates & Forecasts, 1970-2025	6
Table 2 Number of Infant Deaths, Live Births & Infant Death Rates By Race, 1970-2015	7
Figure 2 Comparison of White & Black Infant Death Rates for Michigan Residents, 1970-2015	9
Table 3 Infant Death Rates by County of Residence, 2015 & 2011 - 2015 Average	10
Table 4 Infant Death Rates by Selected Cities of Residence, 2015 & 2011 - 2015 Average	13
Table 5 Infant Deaths by Race for Michigan, Detroit City & Wayne County, 1990- 2015	14
Table 6 Infant Death Rates by Selected Characteristics of Newborn & of Mother, 2015	17
Table 7 Number & Rate of Infant Deaths by Race, Michigan & United States Residents, 1989 - 2015	19
Table 8-A Number of Hispanic Deaths by Selected County & Michigan, 2005 - 2015	20
Table 8-B Number of Arabic Infant Deaths by Selected County & Michigan, 2005 - 2015	21
Table 9 Infant, Hebdomadal, Fetal & Perinatal Death Rates by Specified Race & Ancestry, 2015	22
Table 10 Number of Infant Deaths by Cause of Death, 1980-2015	23
Table 11 Three-year Moving Average Infant Deaths Rates by Cause of Death, 1980-2015	24
Figure 3 Three-year Average Infant Death Rates by Cause, 1980-2015	25
Table 12-A Dollfus Categories, 2014 - 2015	26
Table 12-B Dollfus Subcategories, 2014 - 2015	27
Table 13-A Sudden Unexpected Infant Deaths, 1989 - 2015	28
Table 13-B Accidental Suffocation & Strangulation, 1989 - 2015	29
Table 14 Sudden Unexpected Infant Deaths by Race, 1989 - 2015	30

Three-year Moving Averages by Region

Table 15-A Three-Year Moving Average Infant Death Rates by County, 2004-2015	31
Table 15-B White Three-Year Moving Average Infant Death Rates by County, 2004-2015	34
Table 15-C Black Three-Year Moving Average Infant Death Rates by County, 2004-2015	37
Table 16-A Three-Year Moving Average Infant Death Rates by District, 2004-2015	38

Table Of Contents

Three-year Moving Averages by Region (continued)

Table 16-B White Three-Year Moving Average Infant Death Rates by District, 2004-2015	40
Table 16-C Black Three-Year Moving Average Infant Death Rates by District, 2004-2015	42
Table 16-A Three-Year Moving Average Infant Death Rates by City, 2004-2015	43
Table 17-B White Three-Year Moving Average Infant Death Rates by City, 2004-2015	44
Table 17-C Black Three-Year Moving Average Infant Death Rates by City, 2004-2015	45

Neonatal & Postneonatal Deaths by Region

Table 18-A Neonatal Deaths by Michigan County & Detroit City , 2005-2015	46
Table 18-B Postneonatal Deaths by Michigan County & Detroit City, 2005-2015	50
Table 19-A Neonatal Deaths by Michigan Health Departments, 2005-2015	54
Table 19-B Postneonatal Deaths by Michigan Health Departments, 2005-2015	56

SUMMARY OF 2015 INFANT DEATH STATISTICS

INFANT MORTALITY: For every 1,000 Michigan live births, almost seven infants die before reaching their first birthday. In 2015, 765 infants under the age of one year died, resulting in an infant mortality rate of 6.8 per 1,000 live births. Michigan experienced a significant decline in infant mortality during the early 1990s; but during the 2000s the infant mortality rates remained around 7.9 deaths per 1,000 births. The 2015 rate indicates a drop of the mortality rates to about 7.0 since 2010, from the 7.9 average in the previous decade. (See Table 1.)

Short range forecasts predict within an 80% probability that the mortality rate will gradually decline to about 6.6 by 2020. (See Table 1.)

Both the white and black infant mortality rates remained about the same in recent years, with a persistent racial disparity of about 2.9 black deaths for every 1 white death. In 2005, the white infant death rate was 5.5, and was comparable to the previous ten-year 1996-2005 average of 6.0 deaths per 1,000 white births. The black infant death rate was 17.9 in 2005, and was also comparable to the previous decade average of 17.6 deaths per 1,000 black births. In 2015 the white infant mortality rate was 5.0 per 1,000 live births while the black rate was 14.3 per 1,000 live births. Between 2005-2015 the white infant mortality rate has remained about the same, while the black rate has declined by 25% due to a reduction of infant deaths since 2005. The decline in the Michigan infant mortality rate is largely due to a reduction in black infant deaths. (See Table 2.)

The Michigan infant mortality rate continues to be higher than the last reported national rate. The 2015 infant death rate for the United States is 5.9. (See Table 7.)

LIVE BIRTHS: In 2015, the number of live births decreased to 113,211, from the 2014 live birth count of 114,460. However, the 2011-2015 numbers represent historic lows, and Michigan's birth count has gradually decreased since at least 1990. Nationally, there were 3,977,745 births in the U.S. in 2015, a decline of 7.8% from the record number reported for 2007 and a 0.3% decrease from last year.

CAUSES OF DEATH: In 2015, 29% of infants died due to conditions related to prematurity and 20% died due to birth defects. In addition, 11.6% infants died due to accidents; 9.5% of all infant deaths were due to accidental suffocation in bed. (See Table 12-A.)

Respiratory Distress Syndrome (RDS) declined from about 15 per 1,000 live births in the early eighties to about 5 per 1,000 by the late nineties. Due to surfactant and other therapies, the number of RDS deaths has continued to decline to less than 2 per 1,000 live births in 2015. In the past few years, the SIDS death rate has remained about 3 per 1,000 live births.

CHARACTERISTICS OF NEWBORN: Certain newborns are at higher risk of dying. In 2015 infants born with very low birth weight (less than 1,500 grams) experienced an infant death rate of 227.1 per 1,000 live births compared to a rate of 2.3 for those infants weighing 2,500 grams or more. Multiple birth infants had an infant mortality rate of 26.4 per 1,000 live births compared to the rate of 5.9 for single birth infants. The excess male infant mortality normally reported in other years was present in 2015, with the infant mortality for male infants at 7.4 per 1,000 male live births and females at 6.0 per 1,000 female live births. (See Table 6.)

CHARACTERISTICS OF MOTHER: Infant death rates were the lowest for mothers aged 30-39 years old and highest for mothers aged under 20 years old. Unmarried mothers had infant mortality rates about twice those of married mothers. Women receiving inadequate prenatal care experienced infant mortality rates three times as high as those women receiving adequate prenatal care. Mothers exposed to secondhand smoking while pregnant had an infant death rate of 10.3 per 1,000 live births compared to a rate of 5.8 for mothers who were not exposed to secondhand smoking during pregnancy. (See Table 6.)

COMMUNITY LEVEL DATA: Infant mortality data are available for most communities in Michigan. On the Internet, go to Michigan's community health information site at www.michigan.gov/healthstatistics. Data for local health department districts, counties and major cities and townships can be reached by choosing "Vital Statistics" and then "Infant Mortality" button. Those who do not have access to the Internet may request that community level data be mailed to them from the address below.

Questions regarding Infant Death Statistics should be addressed to:

Lindsey Myers
Division for Vital Records and Health Statistics
Michigan Department of Health & Human Services
333 S. Grand Ave.
Lansing, MI 48933

(517) 335-8715 Voice
(517) 335-8711 FAX

E-mail: MyersL@Michigan.gov

Residence Determined by Reported Residence and Geocoded Residence in the Michigan Death Certificate Registry

Using a process called geocoding, the Division for Vital Records & Health Statistics (DVRHS) has assigned geographic coordinates to most death certificates reported between 2000 and 2014. While the primary purpose of geocoding mortality data is to map patterns of death and to conduct spatial epidemiology, such records can also be used to improve the accuracy of location-related data recorded on the certificate. In mortality datasets, the division has found that the MCD residence is miscoded in certain areas about 10-20% of the time. Often, the person reporting the residence confuses the township with an adjacent city or village. Such errors then alter the calculation of local area statistics; small errors in coding lead to large errors in rates calculated for MCDs with populations below 70,000.

Therefore, all mortality and natality statistics between 2000 and 2014 in division reports of infant deaths were updated and recalculated using the geocoded data. **Birth and death counts and rates are different from data reported prior to 2011.** Statistics for years before 2000 use the residence reported during the course of registering a vital event. When rates calculated from these data differ substantially from Michigan or county rates, one should study the events as part of a trend of geocoded statistics, and consider whether the data reflects jumps indicative of a longterm trend of residence-miscoding.

Table 1
Number of Infant Deaths
Live Births and Infant Death Rates
Michigan Residents, 1970 - 2015


Year	Infant Deaths	Live Births	Infant Death Rate
1970	3,492	171,667	20.3 ±0.7
1971	3,157	162,244	19.5 ±0.7
1972	2,801	146,854	19.1 ±0.7
1973	2,561	141,550	18.1 ±0.7
1974	2,387	137,414	17.4 ±0.7
1975	2,205	133,931	16.5 ±0.7
1976	1,978	131,378	15.1 ±0.7
1977	1,945	138,416	14.1 ±0.6
1978	1,931	138,802	13.9 ±0.6
1979	1,921	144,452	13.3 ±0.6
1980	1,851	145,162	12.8 ±0.6
1981	1,851	140,579	13.2 ±0.6
1982	1,672	137,950	12.1 ±0.6
1983	1,573	133,026	11.8 ±0.6
1984	1,595	135,782	11.7 ±0.6
1985	1,575	138,052	11.4 ±0.6
1986	1,565	137,626	11.4 ±0.6
1987	1,538	140,466	10.9 ±0.5
1988	1,542	139,635	11.0 ±0.5
1989	1,645	148,164	11.1 ±0.5
1990	1,638	153,080	10.7 ±0.5
1991	1,554	149,478	10.4 ±0.5
1992	1,460	143,827	10.2 ±0.5
1993	1,319	139,560	9.5 ±0.5
1994	1,184	137,844	8.6 ±0.5
1995	1,110	134,169	8.3 ±0.5

Table 1
Number of Infant Deaths
Live Births and Infant Death Rates
Michigan Residents, 1970 - 2015
(Continued)

Year	Infant Deaths	Live Births	Infant Death Rate
1996	1,072	133,231	8.0 ±0.5
1997	1,085	133,549	8.1 ±0.5
1998	1,091	133,649	8.2 ±0.5
1999	1,071	133,429	8.0 ±0.5
2000	1,112	136,048	8.2 ±0.5
2001	1,066	133,247	8.0 ±0.5
2002	1,054	129,518	8.1 ±0.5
2003	1,112	130,850	8.5 ±0.5
2004	984	129,710	7.6 ±0.5
2005	1,013	127,518	7.9 ±0.5
2006	940	127,537	7.4 ±0.5
2007	997	125,172	8.0 ±0.5
2008	894	121,231	7.4 ±0.5
2009	881	117,309	7.5 ±0.5
2010	817	114,717	7.1 ±0.5
2011	749	114,159	6.6 ±0.5
2012	783	112,708	6.9 ±0.5
2013	799	113,732	7.0 ±0.5
2014	773	114,460	6.8 ±0.5
2015	765	113,211	6.8 ±0.5

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Source: 1970 - 2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services



Note: Rates are per 1,000 live births. Rates after 2015 are forecasted within an 80% prediction interval indicated by the dotted lines; this means that the future rates fall between the lower and upper bounds of the interval with 80% probability.

Source: 1970- 2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 2
Number of Infant Deaths, Live Births and Infant Death Rates By Race
Michigan Residents, 1970 - 2015

Year	White			Black			Other ¹		
	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate
1970	2,661	144,159	18.5 ±0.7	811	26,513	30.6 ±2.1	14	871	16.1 ±8.4
1971	2,293	134,486	17.1 ±0.7	843	26,755	31.5 ±2.1	17	890	19.1 ±9.0
1972	1,957	120,042	16.3 ±0.7	835	25,760	32.4 ±2.2	8	936	8.5 ±5.9
1973	1,749	115,150	15.2 ±0.7	796	25,252	31.5 ±2.2	11	926	11.9 ±7.0
1974	1,673	112,768	14.8 ±0.7	705	22,836	30.9 ±2.2	7	950	7.4 ±5.4
1975	1,574	110,058	14.3 ±0.7	616	22,692	27.1 ±2.1	8	951	8.4 ±5.8
1976	1,383	108,295	12.8 ±0.7	575	21,778	26.4 ±2.1	16	1,100	14.5 ±7.1
1977	1,373	114,511	12.0 ±0.6	559	22,620	24.7 ±2.0	8	1,059	7.6 ±5.2
1978	1,344	114,652	11.7 ±0.6	566	22,423	25.2 ±2.1	13	1,507	8.6 ±4.7
1979	1,352	119,218	11.3 ±0.6	550	23,346	23.6 ±1.9	12	1,622	7.4 ±4.2
1980	1,246	119,698	10.4 ±0.6	582	23,422	24.8 ±2.0	15	1,825	8.2 ±4.1
1981	1,261	116,596	10.8 ±0.6	564	22,002	25.6 ±2.1	18	1,847	9.7 ±4.5
1982	1,110	114,465	9.7 ±0.6	548	21,574	25.4 ±2.1	13	1,797	7.2 ±3.9
1983	1,048	109,892	9.5 ±0.6	506	21,295	23.8 ±2.0	15	1,721	8.7 ±4.4
1984	1,034	111,574	9.3 ±0.6	541	22,322	24.2 ±2.0	15	1,792	8.4 ±4.2
1985	1,053	114,150	9.2 ±0.6	510	21,935	23.3 ±2.0	11	1,820	6.0 ±3.6
1986	999	112,255	8.9 ±0.5	553	23,366	23.7 ±1.9	12	1,927	6.2 ±3.5
1987	953	113,386	8.4 ±0.5	567	24,975	22.7 ±1.8	17	1,992	8.5 ±4.0
1988	952	112,232	8.5 ±0.5	570	25,217	22.6 ±1.8	19	2,093	9.1 ±4.1
1989	946	115,186	8.2 ±0.5	672	29,839	22.5 ±1.7	23	2,525	9.1 ±3.7
1990	930	118,180	7.9 ±0.5	689	31,842	21.6 ±1.6	18	2,623	6.9 ±3.2
1991	857	114,983	7.5 ±0.5	683	31,543	21.7 ±1.6	13	2,522	5.2 ±2.8
1992	781	111,458	7.0 ±0.5	656	29,656	22.1 ±1.7	23	2,223	10.3 ±4.2
1993	770	108,329	7.1 ±0.5	530	28,249	18.8 ±1.6	17	2,418	7.0 ±3.3
1994	670	106,828	6.3 ±0.5	495	27,129	18.2 ±1.6	19	3,137	6.1 ±2.7
1995	655	105,274	6.2 ±0.5	432	24,914	17.3 ±1.6	19	3,124	6.1 ±2.7
1996	627	104,922	6.0 ±0.5	421	24,073	17.5 ±1.7	21	3,307	6.4 ±2.7
1997	640	104,980	6.1 ±0.5	425	24,201	17.6 ±1.7	17	3,628	4.7 ±2.2
1998	660	105,161	6.3 ±0.5	405	24,179	16.8 ±1.6	23	3,833	6.0 ±2.4
1999	616	104,493	5.9 ±0.5	428	23,850	17.9 ±1.7	19	4,210	4.5 ±2.0

Table 2
Number of Infant Deaths, Live Births and Infant Death Rates By Race
Michigan Residents, 1970 - 2015

Year	White			Black			Other ¹		
	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate
2000	637	106,322	6.0 ±0.5	437	24,069	18.2 ±1.7	30	4,525	6.6 ±2.4
2001	635	104,489	6.1 ±0.5	398	23,494	16.9 ±1.7	25	4,752	5.3 ±2.1
2002	611	101,611	6.0 ±0.5	410	22,248	18.4 ±1.8	21	5,056	4.2 ±1.8
2003	680	102,146	6.7 ±0.5	391	22,380	17.5 ±1.7	37	5,508	6.7 ±2.2
2004	527	100,795	5.2 ±0.4	388	22,484	17.3 ±1.7	61	5,719	10.7 ±2.7
2005	549	99,117	5.5 ±0.5	400	22,365	17.9 ±1.7	62	5,461	11.4 ±2.8
2006	534	98,551	5.4 ±0.5	339	22,873	14.8 ±1.6	60	5,636	10.6 ±2.7
2007	549	93,868	5.8 ±0.5	368	22,314	16.5 ±1.7	74	6,932	10.7 ±2.4
2008	496	91,509	5.4 ±0.5	332	22,695	14.6 ±1.6	61	6,691	9.1 ±2.3
2009	470	87,471	5.4 ±0.5	346	22,292	15.5 ±1.6	46	6,980	6.6 ±1.9
2010	462	84,597	5.5 ±0.5	312	22,046	14.2 ±1.6	38	7,425	5.1 ±1.6
2011	411	84,172	4.9 ±0.5	300	21,950	13.7 ±1.5	34	7,529	4.5 ±1.5
2012	455	82,949	5.5 ±0.5	291	21,623	13.5 ±1.5	35	7,696	4.5 ±1.5
2013	473	83,339	5.7 ±0.5	290	22,055	13.1 ±1.5	35	8,081	4.3 ±1.4
2014	452	84,492	5.3 ±0.5	284	21,282	13.3 ±1.5	36	8,574	4.2 ±1.4
2015	418	82,938	5.0 ±0.5	306	21,457	14.3 ±1.6	40	8,583	4.7 ±1.4


Note: Infant deaths by race of infant; live births used in calculating infant death rates are by race of mother. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Infant deaths of unknown race are not included in this table.

¹Between 2004-2008, multi-racial infants were reported as "other."

Source: 1970 - 2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Figure 2
Comparison of White and Black Infant Death Rates for Michigan Residents, 1970-2015



Note: Rates are per 1,000 live births. The range indicated by the grey band represents a 95% confidence interval; this means that the true rate lies between the lower and upper bounds of the interval with 95% statistical confidence.

Source: 1970-2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2015 and 2011 - 2015 Average

County of Residence	2015			2011 - 2015		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
MICHIGAN	765	113,211	6.8 ±0.5	773.8	113,654.0	6.8 ±0.2
Alcona	-	54	-	0.6	60.0	*
Alger	-	65	-	-	60.0	-
Allegan	10	1,397	7.2 ±4.4	7.8	1,362.6	5.7 ±1.8
Alpena	1	266	*	1.2	267.8	4.5 ±3.6
Antrim	-	205	-	1.0	200.2	*
Arenac	-	137	-	1.8	128.2	14.0 ±9.1
Baraga	1	72	*	0.6	77.4	*
Barry	-	652	-	2.2	627.4	3.5 ±2.1
Bay	6	1,019	5.9 ±4.7	6.4	1,057.8	6.1 ±2.1
Benzie	1	147	*	0.6	150.2	*
Berrien	9	1,726	5.2 ±3.4	14.0	1,829.8	7.7 ±1.8
Branch	3	536	*	3.6	527.4	6.8 ±3.1
Calhoun	14	1,654	8.5 ±4.4	11.8	1,661.4	7.1 ±1.8
Cass	3	499	*	3.6	479.6	7.5 ±3.5
Charlevoix	3	218	*	1.0	237.6	*
Cheboygan	1	223	*	0.2	206.4	*
Chippewa	-	347	-	2.0	358.8	5.6 ±3.4
Clare	1	315	*	1.2	323.6	3.7 ±3.0
Clinton	8	815	9.8 ±6.8	3.8	794.2	4.8 ±2.1
Crawford	2	128	*	0.8	115.0	*
Delta	1	360	*	2.0	375.2	5.3 ±3.3
Dickinson	-	236	-	1.0	249.8	*
Eaton	9	1,237	7.3 ±4.7	7.2	1,182.2	6.1 ±2.0
Emmet	-	314	-	1.0	308.0	*
Genesee	43	4,747	9.1 ±2.7	40.4	4,932.6	8.2 ±1.1
Gladwin	4	277	*	2.0	250.8	8.0 ±4.9
Gogebic	-	131	-	0.2	122.4	*
Gd. Traverse	9	913	9.9 ±6.4	4.0	930.0	4.3 ±1.9
Gratiot	1	402	*	2.2	413.2	5.3 ±3.1
Hillsdale	5	511	*	2.4	539.8	4.4 ±2.5

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2015 and 2011 - 2015 Average

County of Residence	2015			2011 - 2015		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Houghton	-	378	-	1.4	373.0	3.8 ±2.8
Huron	-	311	-	1.0	289.4	*
Ingham	21	3,157	6.7 ±2.8	22.6	3,276.0	6.9 ±1.3
Ionia	3	750	*	3.0	738.0	4.1 ±2.1
Iosco	1	233	*	1.4	232.6	6.0 ±4.4
Iron	-	81	-	0.4	90.6	*
Isabella	5	678	*	3.8	667.0	5.7 ±2.6
Jackson	10	1,729	5.8 ±3.6	10.2	1,786.0	5.7 ±1.6
Kalamazoo	20	3,179	6.3 ±2.7	18.4	3,143.0	5.9 ±1.2
Kalkaska	-	182	-	1.8	173.4	10.4 ±6.7
Kent	56	8,831	6.3 ±1.7	51.2	8,863.0	5.8 ±0.7
Keweenaw	-	16	-	-	18.6	-
Lake	1	111	*	0.4	99.0	*
Lapeer	4	796	*	3.0	823.4	3.6 ±1.8
Leelanau	1	177	*	1.2	170.2	7.1 ±5.6
Lenawee	6	1,014	5.9 ±4.7	7.6	1,059.0	7.2 ±2.3
Livingston	7	1,830	3.8 ±2.8	9.4	1,759.4	5.3 ±1.5
Luce	1	48	*	0.6	52.2	*
Mackinac	2	82	*	0.4	82.6	*
Macomb	60	9,397	6.4 ±1.6	56.0	9,251.2	6.1 ±0.7
Manistee	-	195	-	0.6	183.8	*
Marquette	5	614	*	2.8	636.8	4.4 ±2.3
Mason	2	316	*	1.0	293.0	*
Mecosta	3	431	*	4.2	425.2	9.9 ±4.2
Menominee	1	205	*	0.8	207.4	*
Midland	3	859	*	5.0	862.6	5.8 ±2.3
Missaukee	-	174	-	0.6	179.4	*
Monroe	9	1,411	6.4 ±4.2	7.8	1,520.4	5.1 ±1.6
Montcalm	3	704	*	3.8	719.0	5.3 ±2.4
Montmorency	-	79	-	0.4	66.6	*

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2015 and 2011 - 2015 Average

County of Residence	2015			2011 - 2015		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Muskegon	18	2,079	8.7 ±4.0	17.0	2,106.6	8.1 ±1.7
Newaygo	3	522	*	4.4	549.6	8.0 ±3.3
Oakland	71	13,588	5.2 ±1.2	81.4	13,439.2	6.1 ±0.6
Oceana	2	311	*	2.8	305.4	9.2 ±4.8
Ogemaw	-	167	-	0.8	189.0	*
Ontonagon	-	32	-	0.4	29.8	*
Osceola	2	245	*	1.0	270.4	*
Oscoda	-	103	-	0.6	87.0	*
Otsego	1	260	*	2.8	247.2	11.3 ±5.9
Ottawa	21	3,265	6.4 ±2.7	19.6	3,338.4	5.9 ±1.2
Presque Isle	2	99	*	1.4	92.6	15.1 ±11.1
Roscommon	1	168	*	0.6	173.0	*
Saginaw	19	2,256	8.4 ±3.8	17.8	2,281.0	7.8 ±1.6
St. Clair	9	1,520	5.9 ±3.9	9.6	1,580.0	6.1 ±1.7
St. Joseph	1	792	*	4.8	817.8	5.9 ±2.3
Sanilac	4	440	*	2.6	439.2	5.9 ±3.2
Schoolcraft	-	55	-	-	66.2	-
Shiawasee	4	695	*	2.8	715.0	3.9 ±2.0
Tuscola	4	563	*	2.8	555.2	5.0 ±2.6
Van Buren	6	893	6.7 ±5.4	5.4	910.8	5.9 ±2.2
Washtenaw	14	3,704	3.8 ±2.0	18.8	3,723.6	5.0 ±1.0
Wayne	222	23,472	9.5 ±1.2	225.0	23,456.2	9.6 ±0.6
Wexford	1	407	*	3.4	408.6	8.3 ±3.9
Unknown	-	4	-	-	1.0	-

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2011 - 2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 4
Number of Infant Deaths, Live Births and Infant Death Rates for
Selected Cities of Residence, 2015 and 2011 - 2015 Average**

City of Residence	2015			2011 - 2015		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Ann Arbor	2	1,048	*	4.2	1,099.0	3.8 ±1.6
Battle Creek	9	810	11.1 ±7.2	7.8	841.8	9.3 ±2.9
Dearborn	12	1,661	7.2 ±4.1	11.0	1,656.8	6.6 ±1.7
Dearborn Heights	5	893	*	5.2	804.2	6.5 ±2.5
Detroit	139	9,891	14.1 ±2.3	136.8	10,036.6	13.6 ±1.0
East Lansing	-	-	-	-	-	-
Farmington Hills	4	907	*	5.2	855.2	6.1 ±2.3
Flint	20	1,458	13.7 ±6.0	18.4	1,631.2	11.3 ±2.3
Grand Rapids	23	3,150	7.3 ±3.0	24.2	3,298.4	7.3 ±1.3
Kalamazoo	13	1,039	12.5 ±6.8	8.2	1,139.8	7.2 ±2.2
Kentwood	8	824	9.7 ±6.7	4.8	792.4	6.1 ±2.4
Lansing	14	1,784	7.8 ±4.1	14.6	1,859.0	7.9 ±1.8
Livonia	2	918	*	3.6	874.4	4.1 ±1.9
Midland	2	488	*	2.8	483.8	5.8 ±3.0
Muskegon	8	523	15.3 ±10.5	7.4	596.2	12.4 ±4.0
Novi	1	662	*	2.6	647.8	4.0 ±2.2
Pontiac	18	1,131	15.9 ±7.3	12.6	1,104.8	11.4 ±2.8
Portage	3	586	*	2.6	574.4	4.5 ±2.5
Rochester Hills	2	731	*	3.4	751.8	4.5 ±2.1
Roseville	9	583	15.4 ±10.0	5.2	593.2	8.8 ±3.4
Royal Oak	1	797	*	4.2	761.8	5.5 ±2.4
Saginaw	12	818	14.7 ±8.2	11.2	853.8	13.1 ±3.4
Southfield	5	781	*	7.0	782.0	9.0 ±3.0
St Clair Shores	4	602	*	3.0	580.2	5.2 ±2.6
Sterling Heights	9	1,466	6.1 ±4.0	8.6	1,429.8	6.0 ±1.8
Taylor	5	860	*	7.8	864.4	9.0 ±2.8
Troy	3	838	*	2.8	810.0	3.5 ±1.8
Warren	6	1,678	3.6 ±2.9	11.4	1,660.0	6.9 ±1.8
Westland	6	1,080	5.6 ±4.4	9.4	1,080.8	8.7 ±2.5
Wyoming	8	1,177	6.8 ±4.7	6.2	1,221.0	5.1 ±1.8

Note: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See [Data Quality Considerations](#) for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2011-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2011-2015 Geocoded Michigan Birth Certificate Registries Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2015

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
1990	Total	1,638	10.7 ±0.5	491	20.3 ±1.8	170	10.3 ±1.5	977	8.7 ±0.5
	White	930	7.9 ±0.5	36	8.7 ±2.8	130	8.8 ±1.5	764	7.7 ±0.5
	Black	689	21.6 ±1.6	451	23.0 ±2.1	38	24.8 ±7.8	200	18.7 ±2.6
1991	Total	1,554	10.4 ±0.5	456	19.4 ±1.8	139	8.4 ±1.4	959	8.8 ±0.6
	White	857	7.5 ±0.5	39	10.0 ±3.1	103	7.1 ±1.4	715	7.4 ±0.5
	Black	683	21.7 ±1.6	417	21.6 ±2.1	34	20.3 ±6.7	232	21.9 ±2.8
1992	Total	1,460	10.2 ±0.5	462	21.4 ±1.9	128	8.1 ±1.4	870	8.2 ±0.5
	White	781	7.0 ±0.5	43	12.3 ±3.7	92	6.6 ±1.4	646	6.9 ±0.5
	Black	656	22.1 ±1.7	415	23.3 ±2.2	34	21.8 ±7.2	207	20.1 ±2.7
1993	Total	1,319	9.5 ±0.5	348	17.2 ±1.8	111	7.1 ±1.3	860	8.3 ±0.6
	White	770	7.1 ±0.5	28	9.1 ±3.3	88	6.4 ±1.3	654	7.2 ±0.5
	Black	530	18.8 ±1.6	317	18.7 ±2.0	21	14.4 ±6.1	192	19.4 ±2.7
1994	Total	1,184	8.6 ±0.5	322	16.6 ±1.8	105	6.8 ±1.3	757	7.3 ±0.5
	White	670	6.3 ±0.5	28	9.4 ±3.5	69	5.1 ±1.2	573	6.3 ±0.5
	Black	495	18.2 ±1.6	293	18.2 ±2.1	35	23.7 ±7.7	167	17.5 ±2.6
1995	Total	1,110	8.3 ±0.5	274	15.6 ±1.8	89	5.8 ±1.2	747	7.4 ±0.5
	White	655	6.2 ±0.5	17	6.0 ±2.8	64	4.8 ±1.2	574	6.4 ±0.5
	Black	432	17.3 ±1.6	254	17.6 ±2.1	25	18.2 ±7.1	153	16.8 ±2.6
1996	Total	1,072	8.0 ±0.5	248	15.0 ±1.8	101	6.7 ±1.3	723	7.1 ±0.5
	White	627	6.0 ±0.5	9	3.4 ±2.2	74	5.6 ±1.3	544	6.1 ±0.5
	Black	421	17.5 ±1.7	238	17.5 ±2.2	23	16.2 ±6.6	160	17.7 ±2.7
1997	Total	1,085	8.1 ±0.5	253	15.1 ±1.8	100	6.6 ±1.3	732	7.2 ±0.5
	White	640	6.1 ±0.5	23	8.6 ±3.5	76	5.8 ±1.3	541	6.1 ±0.5
	Black	425	17.6 ±1.7	229	16.7 ±2.1	23	16.5 ±6.7	173	19.1 ±2.8
1998	Total	1,091	8.2 ±0.5	241	14.7 ±1.8	99	6.6 ±1.3	751	7.4 ±0.5
	White	660	6.3 ±0.5	15	5.8 ±2.9	74	5.7 ±1.3	571	6.4 ±0.5
	Black	405	16.8 ±1.6	219	16.1 ±2.1	22	14.9 ±6.2	164	18.0 ±2.7

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2015

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
1999	Total	1,071	8.0 ±0.5	245	15.5 ±1.9	97	6.4 ±1.3	729	7.1 ±0.5
	White	616	5.9 ±0.5	15	5.6 ±2.8	72	5.6 ±1.3	529	5.9 ±0.5
	Black	428	17.9 ±1.7	228	17.8 ±2.3	23	15.1 ±6.1	177	18.6 ±2.7
2000	Total	1,112	8.2 ±0.5	232	14.7 ±1.9	98	6.4 ±1.3	782	7.4 ±0.5
	White	637	6.0 ±0.5	20	7.5 ±3.3	64	5.0 ±1.2	553	6.1 ±0.5
	Black	437	18.2 ±1.7	206	16.1 ±2.2	33	20.2 ±6.8	198	20.5 ±2.8
2001	Total	1,066	8.0 ±0.5	210	13.9 ±1.9	81	5.3 ±1.2	775	7.5 ±0.5
	White	635	6.1 ±0.5	14	5.3 ±2.8	58	4.5 ±1.2	563	6.3 ±0.5
	Black	398	16.9 ±1.7	195	15.9 ±2.2	21	12.9 ±5.5	182	18.9 ±2.7
2002	Total	1,054	8.1 ±0.5	239	16.8 ±2.1	103	6.9 ±1.3	712	7.1 ±0.5
	White	611	6.0 ±0.5	20	8.0 ±3.5	70	5.7 ±1.3	521	6.0 ±0.5
	Black	410	18.4 ±1.8	217	19.1 ±2.5	28	16.9 ±6.2	165	17.9 ±2.7
2003	Total	1,112	8.5 ±0.5	227	16.2 ±2.1	120	8.1 ±1.4	765	7.5 ±0.5
	White	680	6.7 ±0.5	28	11.7 ±4.3	85	7.1 ±1.5	567	6.5 ±0.5
	Black	391	17.5 ±1.7	199	17.9 ±2.5	29	16.9 ±6.1	163	17.1 ±2.6
2004	Total	984	7.6 ±0.5	211	15.6 ±2.1	89	6.1 ±1.3	684	6.7 ±0.5
	White	527	5.2 ±0.4	12	5.0 ±2.8	54	4.6 ±1.2	461	5.3 ±0.5
	Black	388	17.3 ±1.7	193	17.9 ±2.5	30	16.1 ±5.7	165	16.8 ±2.5
2005	Total	1,013	7.9 ±0.5	210	16.0 ±2.1	93	6.5 ±1.3	710	7.1 ±0.5
	White	549	5.5 ±0.5	13	5.7 ±3.1	58	5.1 ±1.3	478	5.6 ±0.5
	Black	400	17.9 ±1.7	188	17.9 ±2.5	34	17.7 ±5.9	178	17.9 ±2.6
2006	Total	940	7.4 ±0.5	175	13.5 ±2.0	98	6.8 ±1.4	667	6.7 ±0.5
	White	534	5.4 ±0.5	9	4.0 ±2.6	61	5.5 ±1.4	464	5.4 ±0.5
	Black	339	14.8 ±1.6	153	14.7 ±2.3	31	14.4 ±5.0	155	15.0 ±2.3

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2015

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
2007	Total	997	8.0 ±0.5	179	14.8 ±2.1	95	6.7 ±1.3	723	7.3 ±0.5
	White	549	5.8 ±0.5	13	7.1 ±3.8	59	5.5 ±1.4	477	5.9 ±0.5
	Black	368	16.5 ±1.7	155	16.1 ±2.5	25	11.4 ±4.5	188	17.9 ±2.5
2008	Total	894	7.4 ±0.5	175	14.8 ±2.2	100	7.2 ±1.4	619	6.5 ±0.5
	White	496	5.4 ±0.5	8	4.7 ±3.3	63	5.9 ±1.5	425	5.4 ±0.5
	Black	332	14.6 ±1.6	159	16.7 ±2.6	27	12.3 ±4.6	146	13.3 ±2.1
2009	Total	881	7.5 ±0.5	166	14.8 ±2.2	82	6.1 ±1.3	633	6.8 ±0.5
	White	470	5.4 ±0.5	6	4.3 ±3.5	54	5.4 ±1.4	410	5.4 ±0.5
	Black	346	15.5 ±1.6	152	16.7 ±2.6	19	7.9 ±3.5	175	16.2 ±2.4
2010	Total	817	7.1 ±0.5	146	13.3 ±2.2	87	6.6 ±1.4	584	6.4 ±0.5
	White	462	5.5 ±0.5	10	9.6 ±5.9	44	4.6 ±1.4	408	5.5 ±0.5
	Black	312	14.2 ±1.6	128	14.3 ±2.5	37	15.0 ±4.8	147	13.8 ±2.2
2011	Total	749	6.6 ±0.5	130	12.6 ±2.1	100	7.5 ±1.5	519	5.7 ±0.5
	White	411	4.9 ±0.5	5	*	58	6.0 ±1.5	348	4.7 ±0.5
	Black	300	13.7 ±1.5	119	14.0 ±2.5	37	14.6 ±4.7	144	13.2 ±2.1
2012	Total	783	6.9 ±0.5	151	15.0 ±2.4	87	6.7 ±1.4	545	6.1 ±0.5
	White	455	5.5 ±0.5	12	15.7 ±8.8	56	5.9 ±1.6	387	5.3 ±0.5
	Black	291	13.5 ±1.5	133	16.0 ±2.7	27	11.2 ±4.2	131	12.0 ±2.0
2013	Total	799	7.0 ±0.5	134	13.3 ±2.2	83	6.1 ±1.3	582	6.5 ±0.5
	White	473	5.7 ±0.5	8	10.0 ±6.9	56	5.9 ±1.5	409	5.6 ±0.5
	Black	290	13.1 ±1.5	120	14.5 ±2.6	22	8.2 ±3.4	148	13.4 ±2.1
2014	Total	773	6.8 ±0.5	130	13.2 ±2.3	88	6.5 ±1.4	555	6.1 ±0.5
	White	452	5.3 ±0.5	8	9.6 ±6.6	55	5.9 ±1.6	389	5.2 ±0.5
	Black	284	13.3 ±1.5	120	14.7 ±2.6	25	9.7 ±3.8	139	13.2 ±2.2
2015	Total	765	6.8 ±0.5	139	14.1 ±2.3	83	6.1 ±1.3	543	6.1 ±0.5
	White	418	5.0 ±0.5	9	10.2 ±6.7	61	6.5 ±1.6	348	4.8 ±0.5
	Black	306	14.3 ±1.6	121	14.9 ±2.6	17	6.4 ±3.1	168	15.7 ±2.4

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Table 6
Number of Infant Deaths, Live Births and Infant Death Rates
by Selected Characteristics of Newborn and of Mother
for Michigan Residents, 2015

Characteristics of Newborn	Infant Deaths	Live Births	Infant Death Rate
Sex of Infant	765	113,211	6.8 ±0.5
Male	428	57,708	7.4 ±0.7
Female	334	55,499	6.0 ±0.6
Birthweight	765	113,211	6.8 ±0.5
1,499 Grams or Less	394	1,735	227.1 ±20.0
1,500 - 2,499 Grams	133	7,876	16.9 ±2.8
2,500 Grams or More	235	103,552	2.3 ±0.3
Plurality	765	113,211	6.8 ±0.5
Single Birth	647	109,001	5.9 ±0.5
Multiple Birth	111	4,204	26.4 ±4.8
Characteristics of Mother	Infant Deaths	Live Births	Infant Death Rate
Age of Mother	765	113,211	6.8 ±0.5
< 20	65	6,412	10.1 ±2.5
20 - 29	438	60,831	7.2 ±0.7
30 - 39	233	43,292	5.4 ±0.7
40 +	22	2,673	8.2 ±3.4
Marital Status	765	113,211	6.8 ±0.5
Married	279	64,825	4.3 ±0.5
Unmarried	479	48,384	9.9 ±0.9
Level of Prenatal Care	765	113,211	6.8 ±0.5
Adequate	357	76,051	4.7 ±0.5
Intermediate	224	26,506	8.5 ±1.1
Inadequate	177	10,639	16.6 ±2.4

Table 6
Number of Infant Deaths, Live Births and Infant Death Rates
by Selected Characteristics of Newborn and of Mother
for Michigan Residents, 2015

Exposure to Tobacco	Infant Deaths	Live Births	Infant Death Rate
Smoked During Pregnancy	765	113,211	6.8 ±0.5
No	574	94,767	6.1 ±0.5
Yes	191	18,444	10.4 ±1.5
Secondhand Smoke During Pregnancy	765	113,211	6.8 ±0.5
No	540	92,321	5.8 ±0.5
Yes	165	16,037	10.3 ±1.6
Smoked or Secondhand Smoke During Pregnancy	765	113,211	6.8 ±0.5
No	465	83,977	5.5 ±0.5
Yes	252	25,208	10.0 ±1.2
History of smoking	765	113,211	6.8 ±0.5
No	534	90,387	5.9 ±0.5
Yes	210	22,150	9.5 ±1.3

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence. Records with selected characteristics unknown are included in total only. Level of prenatal care is according to the Kessner Index which is a classification of prenatal care based on the month of pregnancy in which prenatal care began, the number of prenatal visits and the length of pregnancy (i.e. for shorter pregnancies, fewer prenatal visits constitute adequate care.)

Source: 2015 Michigan Resident Birth-Death Matched File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 7
Number and Rate of Infant Deaths by Race,
Michigan and United States Residents, 1989 - 2015

Year	United States						Michigan					
	All Races		White		Black		All Races		White		Black	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
1989	39,655	9.8	25,794	8.0	12,527	18.6	1,645	11.1	946	8.2	672	22.5
1990	38,351	9.2	24,883	7.6	12,290	18.0	1,638	10.7	930	7.9	689	21.6
1991	36,766	8.9	23,657	7.3	11,994	17.6	1,554	10.4	857	7.5	683	21.7
1992	34,628	8.5	22,164	6.9	11,348	16.8	1,460	10.2	781	7.0	656	22.1
1993	33,466	8.4	21,497	6.8	10,887	16.5	1,319	9.5	770	7.1	530	18.8
1994	31,710	8.0	20,504	6.6	10,072	15.8	1,184	8.6	670	6.3	495	18.2
1995	29,583	7.6	19,490	6.3	9,118	15.1	1,110	8.3	655	6.2	432	17.3
1996	28,487	7.3	18,761	6.1	8,730	14.7	1,072	8.0	627	6.0	421	17.5
1997	28,045	7.2	18,539	6.0	8,496	14.2	1,085	8.1	640	6.1	425	17.6
1998	28,371	7.2	18,561	6.0	8,726	14.3	1,091	8.2	660	6.3	405	16.8
1999	27,953	7.1	18,069	5.8	8,832	14.6	1,071	8.0	616	5.9	428	17.9
2000	28,035	6.9	18,144	5.7	8,771	14.0	1,112	8.2	637	6.0	437	18.2
2001	27,568	6.8	17,955	5.7	8,498	14.0	1,066	8.0	635	6.1	398	16.9
2002	28,034	7.0	18,369	5.8	8,524	14.4	1,054	8.1	611	6.0	410	18.4
2003	28,025	6.9	18,440	5.7	8,402	14.0	1,112	8.5	680	6.7	391	17.5
2004	27,936	6.8	18,231	5.7	8,494	13.3	984	7.6	527	5.2	388	17.3
2005	28,440	6.9	18,514	5.7	8,695	13.7	1,013	7.9	549	5.5	400	17.9
2006	28,527	6.7	18,403	5.6	8,858	13.3	940	7.4	534	5.4	339	14.8
2007	29,138	6.8	18,807	5.6	8,944	13.2	997	8.0	549	5.8	368	16.5
2008	28,059	6.6	18,164	5.6	8,543	12.7	894	7.4	496	5.4	332	14.6
2009	26,412	6.4	16,817	5.3	8,312	12.6	881	7.5	470	5.4	346	15.5
2010	24,548	6.1	15,933	5.2	7,388	11.6	817	7.1	462	5.5	312	14.2
2011	23,910	6.1	15,453	5.1	7,058	11.2	749	6.6	411	4.9	300	13.7
2012	23,654	6.0	15,264	5.1	6,896	10.9	783	6.9	455	5.5	291	13.5
2013	23,440	6.0	15,125	5.0	7,123	11.2	799	7.0	473	5.7	290	13.1
2014	23,215	5.8	14,883	5.1	7,076	10.4	773	6.8	452	5.3	284	13.3
2015	23,455	5.9	14,834	5.0	7,289	10.6	765	6.8	418	5.0	306	14.3

Note: Rates are per 1,000 live births.

Source: 1989 - 2015 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

2007-2012 NCHS, Linked Birth / Infant Death Records 2007-2015 on CDC WONDER On-line Database. Accessed at <http://wonder.cdc.gov/lbd-current.html> on Mar 7, 2017. "Deaths: Final Data for 2006", Vol 57, No. 14, April 2009; and [National Center for Health Statistics, historical data.](#)

Table 8-A
Number of Hispanic and Middle Easterner Infant Deaths by Selected County and Michigan
Michigan Residents, 2005 - 2015

Hispanic Infant Deaths

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	92	94	90	74	74	57	44	74	80	61	77
Hispanic Infant Death Rates	11.0	11.0	10.3	8.5	9.0	7.0	5.6	9.7	10.3	8.0	9.9
Allegan	2	2	-	2	1	-	1	-	3	-	-
Genesee	3	5	5	1	1	1	2	-	-	1	2
Ingham	4	2	3	2	8	1	1	2	4	2	1
Kalamazoo	4	2	4	3	-	4	-	2	2	1	4
Kent	14	27	13	16	6	8	12	10	6	7	13
Lenawee	-	1	2	-	2	1	1	2	5	2	-
Macomb	1	3	6	2	4	-	-	1	3	-	5
Muskegon	1	1	2	2	3	-	1	1	5	-	2
Oakland	13	7	5	9	9	6	7	8	8	4	9
Ottawa	4	4	3	3	4	1	2	4	8	2	4
Saginaw	3	3	8	4	5	2	2	5	4	4	2
Van Buren	7	-	2	2	-	1	-	2	1	2	3
Washtenaw	2	2	-	3	4	2	-	2	1	-	-
Wayne	12	15	19	15	13	15	8	16	17	13	21
All Other Counties	22	20	18	10	14	15	7	19	13	23	11

Table 8-B
Number of Hispanic and Middle Easterner Infant Deaths by Selected County and Michigan
Michigan Residents, 2005 - 2015 (Continued)

Middle Easterner Infant Deaths

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	24	31	32	34	22	24	24	26	36	21	26
Arabic Infant Death Rates	6.3	7.6	8.0	8.5	5.8	5.7	5.9	6.2	7.8	4.4	5.3
Macomb	2	5	3	7	6	5	2	5	6	3	2
Oakland	5	3	4	8	2	6	3	5	7	3	4
Wayne	13	22	23	16	10	10	15	14	17	9	16
All Other Counties	4	1	2	3	4	3	4	2	6	6	4

Note: The increase in hispanic mortality incidence reflects a change in death certificate reporting. Michigan's death certificate was revised in 2003 to comply with a change in the National Standard death certificate and a separate indicator of hispanic ancestry was introduced. In 2005, the hispanic ancestry indicator was used in tabulations for the first time, resulting in the identification of hispanic deaths that might otherwise have been missed. The increase in hispanic death cases between 2004 and 2005 should not be considered as a change in trend.

Caution: Care should be taken drawing inferences from rates based on small numbers of events or small population base. These rates tend to exhibit considerable variation which may negate their usefulness for comparative purpose

Source: 2015 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 9
Infant, Hebdomadal, Fetal and Perinatal Death Rates
by Specified Race and Ancestry
Michigan Residents, 2015

Race/Ancestry	Live Births	Infant Deaths		Hebdomadal Deaths		Fetal Deaths		Perinatal Deaths	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
All Races	113,211	765	6.8 ±0.5	395	3.5 ±0.3	541	4.8 ±0.4	936	8.2 ±0.5
White	82,938	418	5.0 ±0.5	222	2.7 ±0.4	348	4.2 ±0.4	570	6.8 ±0.6
Black	21,457	306	14.3 ±1.6	153	7.1 ±1.1	158	7.3 ±1.1	311	14.4 ±1.6
American Indian	631	6	9.5 ±7.6	3	*	5	*	8	12.6 ±8.7
Asian and Pacific Islander	3,888	12	3.1 ±1.7	4	*	11	2.8 ±1.7	15	3.8 ±1.9
Middle Easterner	4,872	26	5.3 ±2.0	13	2.7 ±1.4	17	3.5 ±1.7	30	6.1 ±2.2
Hispanic	7,768	77	9.9 ±2.2	40	5.1 ±1.6	23	3.0 ±1.2	63	8.1 ±2.0

Note: Hebdomadal deaths are deaths to infants ages zero to six days. Perinatal deaths are hebdomadal plus fetal deaths.

Infant death and hebdomadal rates are per 1,000 live births. Fetal death and perinatal death rates are per 1,000 live births plus fetal deaths.

Rates are calculated using live births by race/ancestry of mother as denominator. Live births are by the race/ancestry of the mother on the birth certificate. Infant and hebdomadal deaths are by the race on the death certificate. Fetal deaths are by the race of the mother on the fetal death certificate. Perinatal deaths are by mother's race/ancestry on the fetal death certificate or the infant's race/ancestry on the death certificate.

Asterisk (*) indicates that data do not meet standards of precision or reliability.

Caution: Care should be taken drawing inferences from rates based on small numbers of events or small population base. These rates tend to exhibit considerable variation which may negate their usefulness for comparative purposes.

Source: 2015 Michigan Resident Infant Death File, 2015 Michigan Fetal Death Registry
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 10
Number of Infant Deaths by Cause of Death,
1980-2015, Michigan Residents

Causes of Death	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015
All Causes	2,280	1,575	1,638	1,110	1,112	1,013	817	749	783	799	773	765
Certain gastrointestinal diseases	7	6	6	6	10	12	9	13	7	5	6	7
Septicemia	8	6	17	2	5	10	4	1	8	2	8	3
Remainder of infectious and parasitic diseases	5	11	18	12	7	6	3	2	2	3	5	4
Meningitis	13	7	11	2	3	3	5	1	2	1	1	-
Pneumonia and Influenza	27	11	26	20	20	6	8	6	5	7	7	5
Congenital anomalies	321	330	300	204	192	197	184	125	127	160	139	152
Newborn affected by maternal complications of pregnancy	72	52	108	47	45	67	40	49	68	52	42	45
Newborn affected by complications of placenta, cord and membranes	50	34	24	33	36	28	26	18	28	30	28	31
Disorders relating to short gestation and unspecified low birthweight	145	150	251	219	262	219	198	188	188	176	174	149
Birth trauma	45	14	16	8	2	1	-	1	-	-	-	-
Intrauterine hypoxia and birth asphyxia	69	43	25	11	24	19	20	7	14	8	7	11
Respiratory distress syndrome	206	176	164	82	65	57	23	14	26	20	19	14
Other respiratory conditions	199	132	120	63	65	52	27	28	27	20	22	17
Infections specific to the perinatal period	35	32	33	23	35	42	27	22	21	22	24	21
Neonatal hemorrhage	38	10	8	13	28	23	12	11	12	9	18	16
All other and ill-define conditions originating in the perinatal period	169	168	73	47	39	46	27	39	34	47	28	39
Sudden infant death syndrome	260	228	257	148	99	47	44	33	15	31	35	33
Accidents and adverse effects	39	35	38	40	55	68	66	83	93	90	89	97
Homicides	7	8	17	12	13	15	13	11	14	5	13	6
All other causes	136	122	126	118	107	95	81	97	92	111	108	115

Note: Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

Source: 1980-2015 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 11
Three-year Moving Average Infant Deaths Rates per 10,000 Live Births, by Cause of Death,
1980-2015, Michigan Residents


Causes of Death	1980 to 1982	1985 to 1987	1990 to 1992	1995 to 1997	2000 to 2002	2005 to 2007	2008 to 2010	2009 to 2011	2010 to 2012	2011 to 2013	2012 to 2014	2013 to 2015
Total Infant Deaths	126.8	112.4	104.2	81.5	81.0	77.6	73.4	70.7	68.8	68.4	69.1	68.5
Certain gastrointestinal diseases	.6	.6	.6	.6	.6	.9	.9	1.0	.8	.7	.5	.5
Septicemia	.4	.5	.9	.4	.5	.6	.6	.4	.4	.3	.5	.4
Remainder of infectious and parasitic diseases	.4	.8	1.1	.6	.5	.5	.3	.2	.2	.2	.3	.4
Meningitis	.9	.4	.4	*	*	.3	.2	.2	.2	*	*	*
Pneumonia and Influenza	1.4	1.1	1.8	1.1	1.1	.5	.6	.6	.6	.5	.6	.6
Congenital anomalies	22.7	21.4	18.1	15.1	13.8	15.1	15.2	14.2	12.8	12.1	12.5	13.2
Newborn affected by maternal complications of pregnancy	5.8	4.6	6.6	2.9	5.7	5.4	4.4	4.4	4.6	5.0	4.8	4.1
Newborn affected by complications of placenta, cord and membranes	3.2	2.6	2.1	2.9	2.4	2.1	2.2	2.0	2.1	2.2	2.5	2.6
Disorders relating to short gestation and unspecified low birthweight	10.6	12.2	18.2	16.0	17.6	17.2	16.7	16.4	16.8	16.2	15.8	14.6
Birth trauma	2.5	.7	.7	.4	.2	*	-	*	*	*	-	-
Intrauterine hypoxia and birth asphyxia	4.5	2.4	1.6	1.2	1.6	.9	1.2	1.0	1.2	.9	.9	.8
Respiratory distress syndrome	14.8	11.2	9.7	6.1	4.5	3.6	2.5	1.7	1.8	1.8	1.9	1.6
Other respiratory conditions	12.2	10.2	7.1	5.1	4.3	3.5	2.8	2.8	2.4	2.2	2.0	1.7
Infections specific to the perinatal period	2.2	2.3	2.2	2.1	3.0	2.9	2.6	2.2	2.0	1.9	2.0	2.0
Neonatal hemorrhage	2.0	.7	.6	.8	1.3	1.3	1.1	1.2	1.0	.9	1.1	1.3
All other and ill-define conditions originating in the perinatal period	13.3	12.1	5.1	3.7	3.1	4.3	3.8	3.7	2.9	3.5	3.2	3.3
Sudden infant death syndrome	16.6	17.1	16.6	10.8	4.3	3.8	3.9	3.5	2.7	2.3	2.4	2.9
Accidents and adverse effects	2.8	2.1	2.4	2.7	4.6	5.7	5.7	6.4	7.1	7.8	8.0	8.1
Homicides	.4	.9	.9	.9	.8	.8	.9	1.0	1.1	.9	.9	.7
All other causes	9.7	8.6	7.6	7.9	8.3	8.3	7.8	7.9	7.9	8.8	9.1	9.8

Note: For purposes of comparing small values, these rates are calculated at deaths per **10,000** live births; in most Michigan Community Health infant death reports, infant deaths are per **1,000**. **Causes of death** are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

An asterisk (*) indicates the rate does not meet standards of reliability or precision. A dash (-) indicates quantity zero.

Source: 1980-2015 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Figure 3
Infant Mortality Rates by Leading Causes of Infant Deaths,
Michigan Residents, 1980-2015



Note: For purposes of comparing small values, these rates are calculated at deaths per **10,000** live births; in most Michigan Community Health infant death reports, infant deaths are per **1,000**.

Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

An asterisk (*) indicates the rate does not meet standards of reliability or precision. A dash (-) indicates quantity zero.

Source: 1980-2015 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

**Table 12-A
Number of Infant Deaths by Cause of Death,
2014-2015, Michigan Residents**

Dollfus Primary Cause of Death Group

Cause of Death Group	Number of Deaths		Percent of Deaths	
	2014	2015	2014	2015
Total	773	765	100.0	100.0
Prematurity & related conditions	247	223	32.0	29.2
Congenital anomalies	139	152	18.0	19.9
Sudden Infant Death Syndrome (SIDS)	35	33	4.5	4.3
Obstetric conditions	47	52	6.1	6.8
Birth asphyxia	7	11	0.9	1.4
Perinatal infections	40	36	5.2	4.7
Other infections	29	25	3.8	3.3
External causes	105	104	13.6	13.6
Other causes	124	129	16.0	16.9

Table 12-A Dollfus Subcategory Cause of Death Group

Dollfus Subcategory	Number of Deaths		Percent of Deaths	
	2014	2015	2014	2015
Total	773	765	100.0%	100.0%
Short gestation & low birthweight	174	149	22.5%	19.5%
Birth trauma, neonatal hemorrhage, & hematological disorders	18	15	2.3%	2.0%
RDS, interstitial emphysema, pulmonary hemorrhage, & atelectasis	35	26	4.5%	3.4%
Chronic respiratory distress syndrome, & respiratory diseases originating in the perinatal period	1	3	0.1%	0.4%
Necrotizing enterocolitis	11	17	1.4%	2.2%
Neonatal cardiac failure & other cardiovascular disorders originating in the perinatal period	8	13	1.0%	1.7%
Congenital malformations, deformations, & chromosomal anomalies, congenital disorders	139	152	18.0%	19.9%
Sudden Infant Death Syndrome	35	33	4.5%	4.3%
Incompetent cervix, premature rupture of membranes	20	26	2.6%	3.4%
Multiple pregnancy	13	12	1.7%	1.6%
Placental previa, separation and placental abnormalities	14	14	1.8%	1.8%
Intrauterine hypoxia, birth asphyxia	7	11	0.9%	1.4%
Streptococcal meningitis	1	0	0.1%	0.0%
Maternal renal and urinary tract diseases, maternal infections, and parasitic diseases	1	0	0.1%	0.0%
Chorioamnionitis	12	13	1.6%	1.7%
Bacterial Sepsis, congenital pneumonia, and other infections specific to the perinatal period	26	23	3.4%	3.0%
Other infectious and parasitic diseases	18	13	2.3%	1.7%
Bacterial haemophilus and pneumococcal meningitis	1	0	0.1%	0.0%
Influenza and pneumonia and other respiratory infections	10	12	1.3%	1.6%
Accidents involving transportation	2	2	0.3%	0.3%
Accidents involving falls	0	0	0.0%	0.0%
Accidental drowning or ingestion causing obstruction to the airways	1	1	0.1%	0.1%
Accidents involving suffocation or strangulation in bed	75	73	9.7%	9.5%
Assault and other intentional injuries	13	6	1.7%	0.8%
Injuries with undetermined intent or unspecified cause	3	1	0.4%	0.1%
Other unintentional injuries	11	21	1.4%	2.7%
Neoplasms, anemias, and other conditions not otherwise specified	124	129	16.0%	16.9%

Note: The categories listed in this table conform to "[Infant Mortality: Development of a Proposed Update to the Dollfus Classification of Infant Deaths](#)" (Nakamura, et. al.), with the exception of certain external causes. Due to the high number of accidental suffocations of infants in bed, the DVRHS lists "Accidents involving suffocation or strangulation in bed" as a separate category.

Source: 2014-2015 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 13-A
Sudden Infant Death Syndrome (SIDS), Accidental Suffocation & Strangulation in Bed (ASSB)
and
Sudden Unexpected Infant Deaths (SUID),
Michigan Residents 1989-2015

Year	Number				Rate Per 10,000 Live Births		
	SIDS	ASSB	Unknown & unspecified cause	SUID	SIDS	ASSB	SUID
1989	254	10	1	265	17.1	0.7	17.9
1990	256	11	2	269	16.7	0.7	17.6
1991	240	13	3	256	16.1	0.9	17.1
1992	243	7	4	254	16.9	0.5	17.7
1993	210	10	10	230	15.0	0.7	16.5
1994	167	10	8	185	12.1	0.7	13.4
1995	148	15	5	168	11.0	1.1	12.5
1996	149	8	5	162	11.2	0.6	12.2
1997	138	9	13	160	10.3	0.7	12.0
1998	148	11	13	172	11.1	0.8	12.9
1999	114	12	9	135	8.5	0.9	10.1
2000	99	28	6	133	7.3	2.1	9.8
2001	96	39	1	136	7.2	2.9	10.2
2002	85	32	10	127	6.6	2.5	9.8
2003	49	33	14	96	3.7	2.5	7.3
2004	51	49	10	110	3.9	3.8	8.5
2005	47	52	10	109	3.7	4.1	8.5
2006	51	52	9	112	4.0	4.1	8.8
2007	46	56	8	110	3.7	4.5	8.8
2008	48	52	2	102	4.0	4.3	8.4
2009	45	63	4	112	3.8	5.4	9.5
2010	44	52	7	103	3.8	4.5	9.0
2011	33	69	13	115	2.9	6.0	10.1
2012	15	75	16	106	1.3	6.7	9.4
2013	31	72	14	117	2.7	6.3	10.3
2014	35	76	14	125	3.1	6.6	10.9
2015	33	73	19	125	2.9	6.5	11.0

Table 13-B
Accidental Suffocation & Strangulation in Bed and Total Accidents,
Michigan Residents, 1989-2015

Year	ASSB	Other Accidents & adverse effects	Total Accidents	ASSB Percent of Accidents
1989	10	37	47	21.3%
1990	11	27	38	28.9%
1991	13	25	38	34.2%
1992	7	22	29	24.1%
1993	10	35	45	22.2%
1994	10	27	37	27.0%
1995	15	25	40	37.5%
1996	8	28	36	22.2%
1997	9	24	33	27.3%
1998	11	24	35	31.4%
1999	12	24	36	33.3%
2000	28	27	55	50.9%
2001	39	18	57	68.4%
2002	32	40	72	44.4%
2003	33	31	64	51.6%
2004	49	21	70	70.0%
2005	52	16	68	76.5%
2006	52	19	71	73.2%
2007	56	21	77	72.7%
2008	52	10	62	83.9%
2009	63	11	74	85.1%
2010	52	13	65	80.0%
2011	69	14	83	83.1%
2012	75	17	92	81.5%
2013	72	18	90	80.0%
2014	76	14	90	84.4%
2015	73	24	97	75.3%

Notes: Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. Before 1999, causes of death were classified according to the following ICD-9 codes: SIDS: 798.0, ASSB: E913.0, Unknown: 799.9. On or after 1999, deaths were classified according to the following ICD-10 codes: SIDS: R95, ASSB: W75 and Unknown: R99.

After 1996, SUID death-scene guidelines and subsequent reporting instructions have shifted how SIDS and ASSB deaths were coded. As a result, some deaths that may have been attributed to SIDS before 1999 were classified as ASSB after 1999.

For further information on sudden unexplained infant deaths, see the [Sudden Unexpected Infant Death Case Registry](#).

Source: 1989 - 2015 Michigan Resident Death Files and Michigan Resident Birth Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 14
Three-year Average Mortality Rates by Race
Sudden Infant Death Syndrome (SIDS), Accidental Suffocation & Strangulation in Bed (ASSB)
and Sudden Unexpected Infant Deaths (SUID),
Michigan Residents 1989-2015

Trend	All			White			Black		
	SIDS	ASSB	SUIDS	SIDS	ASSB	SUIDS	SIDS	ASSB	SUIDS
1989-1991	16.6	0.8	17.5	13.4	0.5	14.1	27.1	1.7	29.0
1990-1992	16.6	0.7	17.5	13.3	0.6	14.1	28.1	1.1	29.3
1991-1993	16.0	0.7	17.1	12.8	0.7	13.8	28.2	0.8	29.5
1992-1994	14.7	0.6	15.9	11.6	0.6	12.7	27.3	0.8	28.7
1993-1995	12.8	0.9	14.2	9.7	0.6	10.9	25.9	1.7	28.3
1994-1996	11.4	0.8	12.7	8.2	0.6	9.3	26.1	1.6	28.1
1995-1997	10.8	0.8	12.2	7.7	0.7	9.0	24.9	1.2	26.6
1996-1998	10.9	0.7	12.3	7.7	0.6	9.0	24.3	1.1	26.4
1997-1999	10.0	0.8	11.7	7.3	0.6	8.5	21.7	1.7	24.9
1998-2000	9.0	1.3	10.9	6.4	0.9	7.7	19.7	3.1	24.1
1999-2001	7.7	2.0	10.0	5.3	1.6	7.2	18.3	3.8	22.8
2000-2002	7.0	2.5	9.9	5.0	1.9	7.3	16.3	5.6	22.5
2001-2003	5.8	2.6	9.1	4.5	2.2	7.2	12.8	5.3	19.4
2002-2004	4.7	2.9	8.5	3.9	2.2	6.9	9.2	7.0	17.9
2003-2005	3.8	3.5	8.1	3.1	2.9	6.7	7.3	6.7	15.6
2004-2006	3.9	4.0	8.6	3.0	3.0	6.7	7.7	8.9	18.0
2005-2007	3.8	4.2	8.7	3.2	3.1	6.7	6.4	10.1	18.2
2006-2008	3.9	4.3	8.7	3.5	2.9	6.7	5.3	10.8	17.4
2007-2009	3.8	4.7	8.9	3.6	3.0	6.8	4.9	11.9	18.1
2008-2010	3.9	4.7	9.0	3.6	3.1	7.0	4.9	11.0	17.0
2009-2011	3.5	5.3	9.5	3.2	3.5	7.1	5.1	12.8	19.5
2010-2012	2.7	5.7	9.5	2.5	3.9	7.3	3.7	13.7	18.6
2011-2013	2.3	6.3	9.9	2.2	4.3	7.6	2.9	15.7	20.6
2012-2014	2.4	5.8	9.4	2.4	3.7	6.9	2.5	15.1	19.2
2013-2015	2.9	5.7	10.0	2.9	4.0	8.0	3.4	13.3	20.0

Notes: Rates are per 10,000 live births. Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. Before 1999, causes of death were classified according to the following ICD-9 codes: SIDS: 798.0, ASSB: E913.0, Unknown: 799.9. On or after 1999, deaths were classified according to the following ICD-10 codes: SIDS: R95, ASSB: W75 and Unknown: R99.

After 1996, SUID death-scene guidelines and subsequent reporting instructions have shifted how SIDS and ASSB deaths were coded. As a result, some deaths that may have been attributed to SIDS before 1999 were classified as ASSB after 1999.

For further information on sudden unexplained infant deaths, see the [Sudden Unexpected Infant Death Case Registry](#).

Source: 1989 - 2015 Michigan Resident Death Files and Michigan Resident Birth Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 15-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	7.6±0.3	7.8±0.3	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.8±0.3
Alcona	*	*	*	*	*	*	*	*	*	*
Alger	-	-	*	*	*	*	-	-	-	-
Allegan	6.3±2.3	8.4±2.6	6.9±2.4	6.7±2.4	4.2±1.9	4.5±2.0	4.2±2.0	5.0±2.2	5.4±2.3	6.8±2.5
Alpena	*	*	6.7±5.3	*	*	7.3±5.8	*	*	*	*
Antrim	*	*	8.7±6.9	*	11.1±8.2	10.9±8.0	10.0±8.0	*	*	*
Arenac	*	*	*	*	*	*	*	18.9±13.9	19.2±14.1	15.7±12.5
Baraga	*	*	*	*	*	-	-	*	*	*
Barry	5.9±3.2	6.5±3.4	7.8±3.7	7.4±3.7	8.9±4.1	7.9±4.0	6.8±3.7	4.9±3.2	4.8±3.1	3.2±2.5
Bay	6.6±2.6	8.0±2.8	7.5±2.8	6.6±2.6	7.9±2.9	8.3±3.0	7.7±3.0	4.9±2.4	5.4±2.6	5.8±2.7
Benzie	*	11.7±8.6	*	*	*	*	*	*	*	*
Berrien	7.3±2.1	9.4±2.4	8.8±2.3	8.9±2.4	7.8±2.2	8.3±2.3	8.1±2.3	8.3±2.4	7.8±2.3	6.9±2.2
Branch	8.7±4.2	5.3±3.3	3.3±2.6	3.5±2.8	4.9±3.4	6.3±3.9	7.8±4.4	7.0±4.1	6.9±4.1	6.1±3.8
Calhoun	9.6±2.6	11.2±2.8	10.8±2.7	11.5±2.9	10.2±2.8	10.7±2.9	8.7±2.6	7.2±2.3	6.0±2.1	6.8±2.3
Cass	8.6±4.5	8.3±4.3	5.9±3.6	7.8±4.2	7.8±4.4	7.3±4.3	6.6±4.3	7.4±4.6	8.5±4.8	7.3±4.3
Charlevoix	*	*	*	*	*	*	*	*	*	*
Cheboygan	*	*	*	*	*	*	-	-	-	*
Chippewa	7.4±4.8	8.6±5.3	5.4±4.3	*	*	5.3±4.2	7.5±5.2	7.5±5.2	5.6±4.5	*
Clare	7.8±5.4	5.6±4.5	6.6±4.9	5.8±4.7	*	*	*	*	*	*
Clinton	3.4±2.2	4.7±2.7	4.8±2.7	4.7±2.8	4.4±2.7	4.0±2.6	3.8±2.5	2.5±2.0	3.8±2.5	6.3±3.2
Crawford	*	16.8±12.3	19.1±13.1	17.5±12.9	*	*	*	*	*	*
Delta	4.9±3.9	4.9±3.9	*	6.1±4.5	*	5.3±4.2	6.4±4.7	5.4±4.3	7.1±4.9	*
Dickinson	7.3±5.9	7.6±6.0	7.6±6.0	*	8.3±6.6	*	*	*	*	*
Eaton	4.4±2.2	4.4±2.2	5.3±2.4	5.3±2.4	5.9±2.5	5.6±2.5	5.4±2.4	6.1±2.6	6.0±2.6	7.0±2.7
Emmet	*	*	*	*	6.3±5.0	*	*	*	*	*
Genesee	10.5±1.5	9.5±1.4	9.3±1.4	9.0±1.4	7.8±1.4	7.8±1.4	7.1±1.3	7.9±1.4	8.0±1.4	8.4±1.5
Gladwin	9.0±6.7	*	*	*	*	*	*	*	*	11.3±7.3
Gogebic	*	*	*	*	*	*	*	-	*	*
Grand Traverse	5.8±2.7	6.1±2.8	7.0±3.0	6.9±3.0	5.6±2.8	3.3±2.1	2.9±2.0	*	3.5±2.2	5.3±2.7
Gratiot	7.4±4.4	7.0±4.3	8.7±4.9	7.1±4.4	8.1±4.7	8.1±4.8	6.3±4.3	4.8±3.8	5.8±4.2	6.6±4.6
Hillsdale	4.0±2.9	6.8±3.8	9.2±4.5	5.9±3.6	4.2±3.1	*	3.7±3.0	4.3±3.2	*	4.8±3.3

Table 15-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Houghton	5.1±4.1	5.0±4.0	7.3±4.8	6.6±4.5	6.5±4.5	*	5.1±4.0	5.3±4.3	5.4±4.3	*
Huron	*	*	*	*	*	6.3±5.1	6.5±5.2	*	*	-
Ingham	7.2±1.6	7.0±1.6	6.6±1.6	7.0±1.6	5.9±1.5	5.6±1.5	5.8±1.5	7.2±1.7	7.1±1.7	7.1±1.7
Ionia	3.5±2.3	4.8±2.7	3.9±2.4	3.3±2.3	3.0±2.2	4.0±2.6	4.0±2.6	4.1±2.6	4.1±2.7	5.0±2.9
Iosco	*	*	*	*	11.7±8.6	12.6±8.6	9.2±7.4	*	*	*
Iron	*	*	*	*	*	*	*	*	*	*
Isabella	8.1±3.7	9.9±4.1	6.2±3.2	5.4±3.1	4.2±2.7	6.8±3.5	7.1±3.7	6.2±3.5	4.0±2.8	4.4±2.9
Jackson	9.0±2.3	9.4±2.4	7.7±2.2	7.1±2.2	6.5±2.1	6.1±2.1	5.3±1.9	6.2±2.1	5.6±2.0	6.2±2.1
Kalamazoo	6.4±1.6	7.2±1.7	7.1±1.7	8.4±1.8	8.6±1.9	7.6±1.8	6.4±1.6	6.0±1.6	6.4±1.6	6.5±1.6
Kalkaska	*	*	-	*	*	*	11.3±9.0	11.6±9.2	17.0±11.0	*
Kent	8.0±1.0	7.6±1.0	7.8±1.0	6.9±1.0	6.8±1.0	6.4±1.0	6.2±0.9	5.6±0.9	5.3±0.9	5.5±0.9
Keweenaw	*	*	*	-	-	-	-	-	-	-
Lake	*	*	*	*	*	*	*	*	*	*
Leapeer	3.3±2.1	4.8±2.5	5.4±2.7	7.4±3.2	8.6±3.6	7.1±3.3	3.6±2.4	2.4±1.9	4.4±2.6	5.3±2.9
Leelanau	*	*	*	*	*	*	*	*	*	*
Lenawee	5.6±2.4	6.7±2.6	8.3±3.0	8.5±3.1	10.8±3.5	9.4±3.3	9.2±3.3	7.2±2.9	8.2±3.1	7.6±3.0
Livingston	3.8±1.5	5.3±1.9	5.5±1.9	6.7±2.2	6.3±2.1	6.7±2.2	5.9±2.1	5.2±2.0	5.4±2.0	5.1±1.9
Luce	*	*	*	*	*	-	-	-	*	*
Mackinac	*	*	*	*	*	*	-	-	-	*
Macomb	5.6±0.8	6.0±0.9	6.1±0.9	7.3±1.0	6.7±1.0	6.5±1.0	5.6±0.9	6.2±0.9	6.0±0.9	6.2±0.9
Manistee	9.8±7.2	*	*	*	*	*	*	*	*	*
Marquette	3.8±2.8	4.6±3.0	4.0±2.8	3.9±2.7	4.1±2.8	5.1±3.1	4.1±2.8	4.1±2.9	*	4.3±3.0
Mason	*	6.1±4.8	7.0±5.2	*	6.4±5.1	*	*	*	*	*
Mecosta	6.5±4.2	4.4±3.5	9.7±5.2	8.2±4.8	9.0±5.1	7.0±4.5	10.2±5.5	11.8±6.0	9.3±5.2	8.7±5.1
Menominee	*	*	*	*	*	*	*	*	*	*
Midland	8.7±3.5	7.0±3.1	5.3±2.8	4.2±2.5	3.9±2.4	4.2±2.5	4.6±2.6	6.6±3.1	7.1±3.3	5.8±2.9
Missaukee	*	*	-	*	*	*	*	*	*	*
Monroe	7.3±2.3	6.8±2.2	6.7±2.2	5.7±2.0	5.7±2.1	4.3±1.8	4.4±1.9	5.8±2.2	5.9±2.2	6.0±2.3
Montcalm	3.7±2.4	2.9±2.1	3.7±2.4	3.9±2.6	5.8±3.1	6.4±3.3	6.8±3.4	6.4±3.3	5.5±3.1	4.6±2.9
Montmorency	*	*	*	-	*	*	*	*	*	-

**Table 15-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015**

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Muskegon	6.7±1.9	6.3±1.8	6.7±1.9	7.1±2.0	7.9±2.1	8.0±2.1	8.7±2.3	8.2±2.2	8.3±2.2	8.3±2.2
Newaygo	6.8±3.7	9.7±4.4	8.4±4.1	8.2±4.1	5.3±3.5	7.8±4.2	7.3±4.1	7.3±4.1	7.2±4.1	7.9±4.3
Oakland	6.6±0.8	6.6±0.8	6.4±0.8	6.2±0.8	6.2±0.8	5.8±0.7	6.2±0.8	6.3±0.8	6.6±0.8	6.0±0.8
Oceana	10.3±5.8	8.7±5.4	6.0±4.5	6.3±4.6	8.3±5.4	5.9±4.7	9.0±5.9	6.5±5.2	13.4±7.5	10.2±6.6
Ogemaw	*	*	*	*	*	*	*	*	*	*
Ontonagon	*	*	*	*	-	-	-	*	*	*
Osceola	8.8±6.1	9.4±6.5	*	*	*	*	*	*	*	*
Oscoda	-	-	*	*	*	*	*	*	*	*
Otsego	*	8.2±6.1	9.3±6.4	8.6±6.3	*	9.0±6.6	11.9±7.7	13.9±8.6	12.6±8.2	9.4±6.9
Ottawa	5.3±1.4	5.2±1.4	5.9±1.5	5.4±1.4	5.7±1.5	5.6±1.5	5.9±1.5	6.5±1.6	5.6±1.5	5.9±1.5
Presque Isle	*	*	*	*	*	*	21.7±17.1	*	*	*
Roscommon	*	*	*	10.9±8.7	*	*	*	*	*	*
Saginaw	8.7±2.0	9.5±2.2	9.4±2.2	10.5±2.3	8.8±2.2	7.9±2.1	6.1±1.8	7.1±2.0	7.7±2.1	8.7±2.2
St. Clair	7.7±2.2	6.6±2.1	5.8±2.0	6.4±2.1	5.6±2.0	6.3±2.2	6.2±2.2	6.3±2.2	6.0±2.2	5.1±2.1
St. Joseph	8.1±3.3	8.6±3.4	10.3±3.7	11.2±4.0	10.6±4.0	8.3±3.5	7.4±3.4	7.3±3.4	7.9±3.5	5.2±2.8
Sanilac	5.8±3.8	7.7±4.3	7.5±4.4	6.8±4.2	*	*	*	4.6±3.6	6.8±4.4	6.8±4.4
Schoolcraft	*	*	*	*	-	-	-	-	-	-
Shiawassee	7.7±3.3	8.8±3.7	5.4±2.9	5.7±3.1	3.2±2.4	3.3±2.4	3.8±2.6	3.2±2.4	4.2±2.7	4.6±2.9
Tuscola	7.7±3.9	6.2±3.5	3.8±2.8	3.9±2.9	6.0±3.7	6.8±4.0	6.8±4.0	4.9±3.4	4.7±3.3	4.7±3.3
Van Buren	7.7±3.1	6.7±2.8	5.5±2.6	6.9±2.9	6.5±2.9	4.5±2.4	5.6±2.8	6.5±3.0	6.2±3.0	5.6±2.8
Washtenaw	6.0±1.3	5.7±1.3	5.2±1.3	5.3±1.3	5.1±1.3	4.9±1.3	4.7±1.3	4.6±1.3	5.9±1.4	5.4±1.4
Wayne	10.6±0.7	10.5±0.7	10.4±0.7	10.4±0.7	10.2±0.7	9.8±0.7	9.9±0.7	9.7±0.7	9.6±0.7	9.3±0.7
Wexford	6.3±4.4	9.2±5.2	6.8±4.4	7.7±4.8	6.3±4.3	6.4±4.4	9.8±5.5	9.8±5.5	10.8±5.8	6.4±4.4

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 15-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	5.4±0.3	5.6±0.3	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3
Alcona	*	*	*	*	*	*	*	*	*	*
Alger	-	-	*	*	*	*	-	-	-	-
Allegan	5.6±2.2	7.6±2.5	5.9±2.3	5.6±2.2	3.4±1.8	4.0±2.0	4.1±2.0	5.0±2.2	5.4±2.3	6.8±2.6
Alpena	*	*	6.9±5.5	*	*	*	*	*	*	*
Antrim	*	*	8.9±7.1	*	11.7±8.6	11.7±8.6	10.7±8.5	*	*	*
Arenac	*	*	*	*	*	*	*	19.3±14.2	19.6±14.3	16.0±12.7
Baraga	*	*	-	-	-	-	-	-	*	*
Barry	5.5±3.1	6.1±3.3	8.0±3.8	7.5±3.8	9.1±4.2	8.0±4.0	6.9±3.7	5.0±3.3	4.9±3.2	3.3±2.6
Bay	6.4±2.6	7.5±2.8	7.1±2.8	6.1±2.6	7.4±2.9	7.9±3.0	7.6±3.0	5.0±2.5	5.1±2.6	5.5±2.7
Benzie	*	10.3±8.2	*	*	*	*	*	*	*	*
Berrien	5.3±2.1	5.9±2.3	5.9±2.3	4.8±2.1	4.0±1.9	3.6±1.8	4.6±2.1	4.9±2.2	4.9±2.2	4.7±2.1
Branch	8.2±4.2	4.9±3.2	3.4±2.7	3.6±2.9	5.0±3.5	6.5±4.0	6.7±4.1	5.2±3.6	5.2±3.6	5.8±3.8
Calhoun	7.3±2.5	8.5±2.6	7.9±2.6	9.3±2.9	7.4±2.6	7.9±2.8	6.4±2.5	5.9±2.4	5.1±2.2	5.2±2.3
Cass	6.6±4.1	6.1±4.0	5.5±3.8	5.6±3.9	5.9±4.1	5.3±3.9	6.8±4.7	7.6±5.0	9.6±5.4	8.1±4.8
Charlevoix	*	*	*	*	*	*	*	*	*	*
Cheboygan	*	*	*	*	*	*	-	-	-	*
Chippewa	6.8±5.4	8.4±6.2	*	*	*	*	*	*	*	*
Clare	6.9±5.1	5.7±4.6	6.8±5.0	6.0±4.8	*	*	*	*	*	*
Clinton	2.8±2.1	3.8±2.4	4.7±2.8	4.5±2.8	4.6±2.8	4.2±2.7	4.1±2.7	2.7±2.2	4.1±2.7	6.0±3.2
Crawford	*	14.7±11.7	17.0±12.5	15.7±12.5	*	*	*	*	-	*
Delta	5.1±4.1	*	*	*	*	*	5.8±4.7	*	6.5±4.8	*
Dickinson	7.5±6.0	7.9±6.3	8.0±6.4	*	*	*	*	*	*	*
Eaton	4.4±2.3	4.8±2.4	5.1±2.5	3.8±2.1	4.2±2.3	3.9±2.2	4.8±2.5	4.8±2.5	4.8±2.5	5.1±2.6
Emmet	*	*	*	*	6.8±5.4	*	*	*	*	*
Genesee	7.3±1.5	6.4±1.4	6.0±1.4	5.4±1.3	5.1±1.3	5.1±1.4	4.8±1.3	5.1±1.4	5.2±1.4	5.6±1.5
Gladwin	9.1±6.7	*	*	*	*	*	*	*	*	11.4±7.4
Gogebic	*	*	*	*	*	*	*	-	-	-
Grand Traverse	5.3±2.7	6.4±2.9	7.4±3.2	7.3±3.2	6.0±3.0	3.5±2.3	3.1±2.2	*	3.7±2.3	5.6±2.8
Gratiot	7.1±4.4	6.7±4.3	7.7±4.8	6.0±4.2	7.0±4.5	7.8±4.8	5.8±4.3	*	5.2±4.2	6.2±4.6
Hillsdale	4.0±3.0	7.0±4.0	9.6±4.7	6.2±3.8	4.3±3.2	*	3.8±3.1	4.4±3.2	*	4.9±3.4

Table 15-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Houghton	5.4±4.3	5.3±4.2	7.0±4.8	6.1±4.5	6.0±4.4	*	5.4±4.3	5.7±4.6	5.7±4.6	*
Huron	*	*	*	*	*	6.5±5.2	6.7±5.3	*	*	-
Ingham	4.4±1.4	4.6±1.5	4.1±1.4	4.7±1.5	3.6±1.4	4.6±1.6	5.3±1.7	7.5±2.1	7.1±2.0	6.4±1.9
Ionia	3.6±2.3	4.9±2.8	4.1±2.5	3.4±2.4	3.1±2.3	3.6±2.5	3.7±2.6	3.7±2.6	4.2±2.8	5.1±3.0
Iosco	*	*	*	*	12.2±9.0	13.1±9.0	9.6±7.6	*	*	*
Iron	*	*	*	*	*	*	*	*	*	*
Isabella	8.2±4.0	8.3±4.0	5.0±3.1	3.1±2.5	*	5.6±3.4	6.5±3.8	5.9±3.7	3.5±2.8	4.0±3.0
Jackson	6.2±2.1	6.8±2.2	6.4±2.2	6.2±2.2	5.7±2.1	5.5±2.1	4.1±1.8	4.6±2.0	3.9±1.8	5.1±2.1
Kalamazoo	4.6±1.5	5.0±1.6	4.8±1.6	5.5±1.7	5.7±1.7	4.8±1.6	3.9±1.4	3.9±1.4	4.7±1.6	4.7±1.6
Kalkaska	*	*	-	*	*	*	11.7±9.3	11.9±9.5	17.4±11.3	*
Kent	5.4±0.9	5.2±0.9	5.6±1.0	5.0±0.9	5.0±0.9	4.4±0.9	4.4±0.9	4.1±0.9	4.5±0.9	4.9±0.9
Keweenaw	*	*	*	-	-	-	-	-	-	-
Lake	*	*	*	*	*	*	*	*	*	*
Lapeer	3.4±2.1	4.9±2.6	5.5±2.8	7.3±3.3	7.8±3.5	6.3±3.2	3.0±2.2	2.6±2.1	4.7±2.8	5.6±3.0
Leelanau	*	*	*	*	*	*	*	*	*	*
Lenawee	5.6±2.4	6.8±2.8	9.0±3.3	8.4±3.3	10.6±3.7	8.7±3.4	9.6±3.6	7.6±3.2	8.4±3.3	7.7±3.2
Livingston	3.7±1.6	5.1±1.8	5.3±1.9	6.6±2.2	6.1±2.2	6.6±2.2	5.8±2.1	5.0±2.0	5.1±2.0	4.7±1.9
Luce	*	*	*	*	*	-	-	-	*	*
Mackinac	*	*	*	*	*	*	-	-	-	*
Macomb	4.8±0.8	5.1±0.9	5.2±0.9	6.2±1.0	5.9±1.0	5.6±1.0	4.8±0.9	5.5±1.0	5.3±1.0	5.4±1.0
Manistee	10.3±7.6	*	*	*	*	*	*	*	*	*
Marquette	4.0±2.9	4.9±3.2	4.2±2.9	3.6±2.7	3.7±2.8	4.8±3.1	4.2±2.9	4.3±3.0	*	4.5±3.1
Mason	*	*	6.2±4.9	*	6.6±5.3	*	*	*	*	*
Mecosta	6.1±4.2	*	9.4±5.3	7.9±4.9	8.8±5.2	6.6±4.5	10.0±5.6	11.7±6.1	9.9±5.6	9.2±5.4
Menominee	*	*	*	*	*	*	*	*	*	*
Midland	8.3±3.5	7.3±3.3	5.6±2.9	4.0±2.5	3.7±2.4	4.0±2.5	4.9±2.8	5.8±3.0	6.3±3.2	4.9±2.8
Missaukee	*	*	-	*	*	*	*	*	*	*
Monroe	6.9±2.3	6.8±2.4	6.4±2.3	4.8±2.0	4.7±1.9	3.5±1.7	3.6±1.8	5.1±2.1	5.3±2.2	6.0±2.3
Montcalm	3.7±2.4	3.0±2.2	3.8±2.5	4.0±2.6	5.9±3.2	6.1±3.3	6.5±3.4	6.0±3.3	5.6±3.2	4.7±2.9
Montmorency	*	*	*	-	*	*	*	*	*	-

**Table 15-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015**

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Muskegon	4.5±1.8	5.0±1.9	5.7±2.0	6.4±2.2	6.5±2.2	6.3±2.2	7.1±2.4	6.9±2.3	6.0±2.2	4.8±1.9
Newaygo	5.9±3.5	8.9±4.2	8.0±4.0	8.4±4.2	5.4±3.5	7.4±4.2	6.9±4.1	7.0±4.1	7.6±4.3	8.4±4.6
Oakland	4.9±0.7	4.9±0.8	4.9±0.8	4.8±0.8	5.3±0.8	5.1±0.8	5.7±0.9	5.6±0.9	5.4±0.8	4.5±0.8
Oceana	9.6±5.6	7.4±5.1	*	*	5.9±4.7	*	9.6±6.3	7.1±5.6	14.7±8.3	9.9±6.8
Ogemaw	*	*	*	*	*	*	*	*	*	*
Ontonagon	*	*	*	*	-	-	-	*	*	*
Osceola	8.9±6.2	9.5±6.6	*	*	*	*	*	*	*	*
Oscoda	-	-	*	*	*	*	*	*	*	*
Otsego	*	8.3±6.1	9.4±6.5	8.6±6.4	*	9.2±6.8	12.2±7.9	14.4±8.9	11.5±8.0	8.3±6.6
Ottawa	5.0±1.4	5.0±1.4	5.3±1.4	4.6±1.4	4.9±1.4	5.3±1.5	5.6±1.5	6.2±1.6	5.2±1.4	5.5±1.5
Presque Isle	*	*	*	*	*	*	22.1±17.5	*	*	*
Roscommon	*	*	*	11.2±8.9	*	*	*	*	*	*
Saginaw	5.0±1.8	5.7±2.0	6.6±2.2	6.4±2.2	4.7±1.9	2.9±1.5	2.9±1.5	4.2±1.8	6.2±2.2	6.1±2.2
St. Clair	7.0±2.1	5.8±2.0	5.0±1.9	6.2±2.1	5.4±2.1	6.3±2.3	6.2±2.3	6.3±2.3	5.9±2.3	4.3±1.9
St. Joseph	7.0±3.1	7.6±3.2	9.3±3.6	10.2±3.9	9.9±3.9	7.9±3.6	6.9±3.4	6.4±3.2	6.6±3.3	4.3±2.7
Sanilac	5.8±3.8	7.2±4.2	7.0±4.3	6.3±4.1	*	*	*	4.6±3.7	6.9±4.5	6.2±4.3
Schoolcraft	*	*	*	*	-	-	-	-	-	-
Shiawassee	7.8±3.4	9.0±3.8	5.5±3.0	5.9±3.2	3.3±2.4	3.4±2.5	3.9±2.7	3.3±2.5	4.3±2.8	4.3±2.8
Tuscola	7.8±3.9	6.3±3.6	3.9±2.9	4.0±3.0	6.1±3.8	6.9±4.1	7.0±4.1	5.0±3.5	4.9±3.4	4.8±3.3
Van Buren	4.9±2.6	3.6±2.2	4.8±2.6	6.1±3.0	6.0±3.0	4.1±2.5	5.4±2.9	6.2±3.1	5.8±3.0	4.5±2.7
Washtenaw	4.5±1.4	3.5±1.2	3.5±1.3	4.3±1.4	4.4±1.5	4.5±1.5	3.9±1.4	4.2±1.5	5.4±1.7	4.8±1.6
Wayne	5.0±0.7	5.4±0.7	5.6±0.7	5.6±0.8	5.4±0.8	5.5±0.8	5.9±0.9	6.3±0.9	6.4±0.9	6.4±0.9
Wexford	6.5±4.5	9.5±5.3	7.0±4.6	8.0±5.0	6.5±4.5	6.6±4.5	10.1±5.7	9.3±5.4	10.3±5.8	5.8±4.3

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 15-C
Black Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2004-2015**

County	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	16.6±1.0	16.4±1.0	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.6±0.9
Berrien	11.9±5.5	16.9±6.5	14.9±6.0	16.7±6.4	16.0±6.2	19.5±6.9	18.6±6.9	17.3±6.7	16.5±6.7	13.3±6.1
Calhoun	20.8±9.8	25.3±10.7	24.2±10.5	20.5±9.6	22.1±10.4	25.4±11.3	20.2±10.1	13.8±8.1	9.9±6.8	14.6±8.2
Cass	*	*	*	*	*	*	*	*	-	-
Genesee	18.6±3.8	17.1±3.6	17.1±3.6	17.9±3.7	14.6±3.4	14.2±3.4	12.4±3.3	14.2±3.5	14.6±3.6	15.6±3.7
Ingham	19.8±6.8	19.4±6.8	17.8±6.3	16.5±6.0	16.7±5.9	13.1±5.3	11.4±5.0	10.1±4.7	10.1±4.7	12.9±5.2
Jackson	28.9±13.5	28.7±13.4	16.8±10.3	15.6±10.1	13.2±9.1	11.6±8.6	13.0±9.0	18.5±10.8	16.2±10.0	15.0±9.7
Kalamazoo	16.0±6.1	16.8±6.2	16.8±6.2	19.2±6.5	19.5±6.5	18.3±6.4	17.6±6.3	15.5±5.9	14.2±5.7	15.3±5.9
Kent	18.0±4.4	16.8±4.2	14.4±3.8	13.4±3.7	13.9±3.8	16.6±4.2	14.6±3.9	12.9±3.6	9.9±3.2	10.4±3.3
Macomb	12.7±4.5	13.5±4.3	13.4±4.0	15.1±4.0	13.8±3.8	13.1±3.6	11.4±3.2	10.8±3.0	9.6±2.8	10.6±2.9
Muskegon	13.4±6.0	9.8±5.1	9.0±4.9	8.2±4.6	13.2±5.9	14.9±6.3	16.4±6.8	12.8±6.0	14.9±6.7	19.7±7.7
Oakland	16.7±3.3	17.4±3.4	15.4±3.1	14.7±3.0	11.9±2.7	10.8±2.5	10.6±2.4	11.1±2.5	13.7±2.8	14.6±2.9
Saginaw	18.4±5.7	19.0±5.8	15.3±5.2	19.2±5.8	17.4±5.6	19.2±6.1	13.4±5.1	14.3±5.2	12.2±5.0	15.9±5.7
St. Clair	*	*	*	*	*	*	*	*	*	*
Van Buren	*	*	*	*	*	*	*	*	*	*
Washtenaw	12.1±4.6	13.4±4.9	13.2±4.9	11.6±4.6	10.5±4.4	8.7±4.0	8.9±4.1	8.3±4.1	10.9±4.7	12.1±5.2
Wayne	16.7±1.3	15.9±1.3	15.3±1.3	15.3±1.3	15.1±1.3	14.5±1.3	14.5±1.3	14.0±1.3	13.8±1.3	13.1±1.2

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 16-A
Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2004-2015

Health Department	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	7.6±0.3	7.8±0.3	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.8±0.3
Allegan County	6.3±2.3	8.4±2.6	6.9±2.4	6.7±2.4	4.2±1.9	4.5±2.0	4.2±2.0	5.0±2.2	5.4±2.3	6.8±2.5
Barry-Eaton District	5.0±1.8	5.2±1.9	6.2±2.0	6.0±2.0	7.0±2.2	6.4±2.1	5.9±2.0	5.7±2.0	5.6±2.0	5.7±2.0
Bay County	6.6±2.6	8.0±2.8	7.5±2.8	6.6±2.6	7.9±2.9	8.3±3.0	7.7±3.0	4.9±2.4	5.4±2.6	5.8±2.7
Benzie-Leelanau District	7.0±4.8	7.7±5.0	*	*	*	*	*	*	6.2±5.0	8.2±5.7
Berrien County	7.3±2.1	9.4±2.4	8.8±2.3	8.9±2.4	7.8±2.2	8.3±2.3	8.1±2.3	8.3±2.4	7.8±2.3	6.9±2.2
Branch-Hillsdale-St. Joseph	7.1±2.0	7.2±2.1	8.0±2.2	7.5±2.2	7.2±2.2	6.3±2.0	6.4±2.1	6.3±2.1	6.2±2.0	5.4±1.9
Calhoun County	9.6±2.6	11.2±2.8	10.8±2.7	11.5±2.9	10.2±2.8	10.7±2.9	8.7±2.6	7.2±2.3	6.0±2.1	6.8±2.3
Central Michigan District	7.9±2.2	7.9±2.2	5.6±1.9	5.7±1.9	4.6±1.8	6.3±2.1	5.6±2.0	5.6±2.0	4.8±1.9	5.9±2.0
Chippewa County	7.4±4.8	8.6±5.3	5.4±4.3	*	*	5.3±4.2	7.5±5.2	7.5±5.2	5.6±4.5	*
City Of Detroit	15.0±1.2	14.7±1.2	14.3±1.2	14.8±1.3	14.4±1.3	13.6±1.3	13.6±1.3	13.6±1.3	13.9±1.3	13.5±1.3
Dickinson-Iron District	6.5±4.8	6.6±4.9	6.5±4.8	5.6±4.5	8.7±5.7	8.0±5.5	*	*	5.8±4.6	5.9±4.7
District #10	7.1±1.7	7.8±1.8	6.8±1.7	6.7±1.7	6.7±1.7	6.6±1.7	7.6±1.9	7.0±1.8	8.2±1.9	7.1±1.8
District #2	4.5±3.1	4.7±3.3	5.9±3.6	8.4±4.4	9.8±4.8	9.1±4.6	8.9±4.5	6.5±3.8	7.0±3.9	4.1±3.1
District #4	4.4±2.8	5.3±3.0	6.3±3.3	6.1±3.3	6.1±3.4	6.8±3.7	6.5±3.7	4.9±3.2	4.8±3.1	4.2±2.9
Genesee County	10.5±1.5	9.5±1.4	9.3±1.4	9.0±1.4	7.8±1.4	7.8±1.4	7.1±1.3	7.9±1.4	8.0±1.4	8.4±1.5
Grand Traverse County	5.8±2.7	6.1±2.8	7.0±3.0	6.9±3.0	5.6±2.8	3.3±2.1	2.9±2.0	*	3.5±2.2	5.3±2.7
Huron County	*	*	*	*	*	6.3±5.1	6.5±5.2	*	*	-
Ingham County	7.2±1.6	7.0±1.6	6.6±1.6	7.0±1.6	5.9±1.5	5.6±1.5	5.8±1.5	7.2±1.7	7.1±1.7	7.1±1.7
Ionia County	3.5±2.3	4.8±2.7	3.9±2.4	3.3±2.3	3.0±2.2	4.0±2.6	4.0±2.6	4.1±2.6	4.1±2.7	5.0±2.9
Jackson County	9.0±2.3	9.4±2.4	7.7±2.2	7.1±2.2	6.5±2.1	6.1±2.1	5.3±1.9	6.2±2.1	5.6±2.0	6.2±2.1
Kalamazoo County	6.4±1.6	7.2±1.7	7.1±1.7	8.4±1.8	8.6±1.9	7.6±1.8	6.4±1.6	6.0±1.6	6.4±1.6	6.5±1.6
Kent County	8.0±1.0	7.6±1.0	7.8±1.0	6.9±1.0	6.8±1.0	6.4±1.0	6.2±0.9	5.6±0.9	5.3±0.9	5.5±0.9
Lapeer County	3.3±2.1	4.8±2.5	5.4±2.7	7.4±3.2	8.6±3.6	7.1±3.3	3.6±2.4	2.4±1.9	4.4±2.6	5.3±2.9
Lenawee County	5.6±2.4	6.7±2.6	8.3±3.0	8.5±3.1	10.8±3.5	9.4±3.3	9.2±3.3	7.2±2.9	8.2±3.1	7.6±3.0
Livingston County	3.8±1.5	5.3±1.9	5.5±1.9	6.7±2.2	6.3±2.1	6.7±2.2	5.9±2.1	5.2±2.0	5.4±2.0	5.1±1.9
Luce-Mackinac-Alger-Schoolcraft	9.6±6.2	8.5±5.9	13.2±7.4	11.2±6.9	10.4±6.8	*	-	-	*	*
Macomb County	5.6±0.8	6.0±0.9	6.1±0.9	7.3±1.0	6.7±1.0	6.5±1.0	5.6±0.9	6.2±0.9	6.0±0.9	6.2±0.9
Marquette County	3.8±2.8	4.6±3.0	4.0±2.8	3.9±2.7	4.1±2.8	5.1±3.1	4.1±2.8	4.1±2.9	*	4.3±3.0
Midland County	8.7±3.5	7.0±3.1	5.3±2.8	4.2±2.5	3.9±2.4	4.2±2.5	4.6±2.6	6.6±3.1	7.1±3.3	5.8±2.9

Table 16-A
Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2004-2015

Health Department	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Mid-Michigan District	4.4±1.6	4.5±1.6	5.3±1.8	5.0±1.8	5.8±1.9	5.9±2.0	5.5±1.9	4.4±1.7	4.8±1.8	5.7±2.0
Monroe County	7.3±2.3	6.8±2.2	6.7±2.2	5.7±2.0	5.7±2.1	4.3±1.8	4.4±1.9	5.8±2.2	5.9±2.2	6.0±2.3
Muskegon County	6.7±1.9	6.3±1.8	6.7±1.9	7.1±2.0	7.9±2.1	8.0±2.1	8.7±2.3	8.2±2.2	8.3±2.2	8.3±2.2
Northwest Michigan Community Health	4.6±2.2	5.4±2.4	6.1±2.6	6.5±2.8	6.4±2.8	6.8±2.9	6.9±2.9	6.7±2.9	6.2±2.8	5.8±2.7
Oakland County	6.6±0.8	6.6±0.8	6.4±0.8	6.2±0.8	6.2±0.8	5.8±0.7	6.2±0.8	6.3±0.8	6.6±0.8	6.0±0.8
Ottawa County	5.3±1.4	5.2±1.4	5.9±1.5	5.4±1.4	5.7±1.5	5.6±1.5	5.9±1.5	6.5±1.6	5.6±1.5	5.9±1.5
Delta & Menominee	4.7±3.1	4.6±3.0	4.3±3.0	6.0±3.5	4.6±3.2	4.6±3.2	5.4±3.5	4.7±3.2	5.7±3.5	4.0±2.9
Saginaw County	8.7±2.0	9.5±2.2	9.4±2.2	10.5±2.3	8.8±2.2	7.9±2.1	6.1±1.8	7.1±2.0	7.7±2.1	8.7±2.2
Sanilac County	5.8±3.8	7.7±4.3	7.5±4.4	6.8±4.2	*	*	*	4.6±3.6	6.8±4.4	6.8±4.4
Shiawassee County	7.7±3.3	8.8±3.7	5.4±2.9	5.7±3.1	3.2±2.4	3.3±2.4	3.8±2.6	3.2±2.4	4.2±2.7	4.6±2.9
St Clair County	7.7±2.2	6.6±2.1	5.8±2.0	6.4±2.1	5.6±2.0	6.3±2.2	6.2±2.2	6.3±2.2	6.0±2.2	5.1±2.1
Tuscola County	7.7±3.9	6.2±3.5	3.8±2.8	3.9±2.9	6.0±3.7	6.8±4.0	6.8±4.0	4.9±3.4	4.7±3.3	4.7±3.3
Van Buren-Cass County	8.0±2.5	7.2±2.4	5.6±2.1	7.2±2.4	6.9±2.4	5.5±2.2	6.0±2.3	6.8±2.5	7.0±2.5	6.2±2.4
Washtenaw County	6.0±1.3	5.7±1.3	5.2±1.3	5.3±1.3	5.1±1.3	4.9±1.3	4.7±1.3	4.6±1.3	5.9±1.4	5.4±1.4
Wayne County	6.5±0.8	6.7±0.8	6.9±0.8	6.7±0.8	6.6±0.8	6.7±0.8	6.9±0.8	6.8±0.8	6.4±0.8	6.2±0.8
Western Upper Peninsula District	5.7±3.2	6.6±3.4	7.5±3.7	5.2±3.1	5.3±3.1	3.5±2.6	4.1±2.8	4.3±2.9	6.0±3.5	5.4±3.4

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 16-B
White Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2004-2015**

Health Department	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	5.4±0.3	5.6±0.3	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3
Allegan County	5.6±2.2	7.6±2.5	5.9±2.3	5.6±2.2	3.4±1.8	4.0±2.0	4.1±2.0	5.0±2.2	5.4±2.3	6.8±2.6
Barry-Eaton District	4.8±1.9	5.3±2.0	6.3±2.1	5.2±2.0	6.1±2.1	5.5±2.1	5.6±2.1	4.9±2.0	4.9±2.0	4.4±1.9
Bay County	6.4±2.6	7.5±2.8	7.1±2.8	6.1±2.6	7.4±2.9	7.9±3.0	7.6±3.0	5.0±2.5	5.1±2.6	5.5±2.7
Benzie-Leelanau District	6.5±4.8	7.3±5.1	*	*	*	*	*	*	*	6.8±5.4
Berrien County	5.3±2.1	5.9±2.3	5.9±2.3	4.8±2.1	4.0±1.9	3.6±1.8	4.6±2.1	4.9±2.2	4.9±2.2	4.7±2.1
Branch-Hillsdale-St. Joseph	6.5±2.0	6.7±2.0	7.7±2.2	7.2±2.2	6.9±2.2	6.2±2.1	6.0±2.1	5.5±2.0	5.2±1.9	4.9±1.8
Calhoun County	7.3±2.5	8.5±2.6	7.9±2.6	9.3±2.9	7.4±2.6	7.9±2.8	6.4±2.5	5.9±2.4	5.1±2.2	5.2±2.3
Central Michigan District	7.9±2.3	7.3±2.2	5.2±1.9	4.8±1.8	4.0±1.7	5.8±2.1	5.3±2.0	5.6±2.1	4.7±1.9	5.7±2.1
Chippewa County	6.8±5.4	8.4±6.2	*	*	*	*	*	*	*	*
City Of Detroit	4.9±1.6	5.5±1.8	5.2±1.9	5.5±2.1	5.8±2.3	6.5±2.8	10.3±3.9	10.5±4.1	11.7±4.3	9.9±3.9
Dickinson-Iron District	6.6±4.9	6.9±5.1	6.9±5.1	*	8.2±5.6	7.2±5.3	*	*	5.9±4.7	6.1±4.9
District #10	6.8±1.7	7.4±1.8	6.4±1.6	6.3±1.7	6.4±1.7	6.3±1.7	7.7±1.9	6.8±1.8	8.2±2.0	7.0±1.9
District #2	4.7±3.2	4.2±3.1	5.5±3.6	8.0±4.3	10.1±4.9	9.3±4.7	9.2±4.6	6.7±3.9	7.2±4.0	4.2±3.1
District #4	4.1±2.7	5.5±3.1	6.6±3.4	6.4±3.4	6.3±3.6	6.5±3.7	6.2±3.7	4.5±3.1	4.4±3.1	3.8±2.8
Genesee County	7.3±1.5	6.4±1.4	6.0±1.4	5.4±1.3	5.1±1.3	5.1±1.4	4.8±1.3	5.1±1.4	5.2±1.4	5.6±1.5
Grand Traverse County	5.3±2.7	6.4±2.9	7.4±3.2	7.3±3.2	6.0±3.0	3.5±2.3	3.1±2.2	*	3.7±2.3	5.6±2.8
Huron County	*	*	*	*	*	6.5±5.2	6.7±5.3	*	*	-
Ingham County	4.4±1.4	4.6±1.5	4.1±1.4	4.7±1.5	3.6±1.4	4.6±1.6	5.3±1.7	7.5±2.1	7.1±2.0	6.4±1.9
Ionia County	3.6±2.3	4.9±2.8	4.1±2.5	3.4±2.4	3.1±2.3	3.6±2.5	3.7±2.6	3.7±2.6	4.2±2.8	5.1±3.0
Jackson County	6.2±2.1	6.8±2.2	6.4±2.2	6.2±2.2	5.7±2.1	5.5±2.1	4.1±1.8	4.6±2.0	3.9±1.8	5.1±2.1
Kalamazoo County	4.6±1.5	5.0±1.6	4.8±1.6	5.5±1.7	5.7±1.7	4.8±1.6	3.9±1.4	3.9±1.4	4.7±1.6	4.7±1.6
Kent County	5.4±0.9	5.2±0.9	5.6±1.0	5.0±0.9	5.0±0.9	4.4±0.9	4.4±0.9	4.1±0.9	4.5±0.9	4.9±0.9
Lapeer County	3.4±2.1	4.9±2.6	5.5±2.8	7.3±3.3	7.8±3.5	6.3±3.2	3.0±2.2	2.6±2.1	4.7±2.8	5.6±3.0
Lenawee County	5.6±2.4	6.8±2.8	9.0±3.3	8.4±3.3	10.6±3.7	8.7±3.4	9.6±3.6	7.6±3.2	8.4±3.3	7.7±3.2
Livingston County	3.7±1.6	5.1±1.8	5.3±1.9	6.6±2.2	6.1±2.2	6.6±2.2	5.8±2.1	5.0±2.0	5.1±2.0	4.7±1.9
Luce-Mackinac-Alger-Schoolcraft	8.8±6.5	8.8±6.5	11.7±7.6	10.5±7.2	9.5±7.0	*	-	-	*	*
Macomb County	4.8±0.8	5.1±0.9	5.2±0.9	6.2±1.0	5.9±1.0	5.6±1.0	4.8±0.9	5.5±1.0	5.3±1.0	5.4±1.0
Marquette County	4.0±2.9	4.9±3.2	4.2±2.9	3.6±2.7	3.7±2.8	4.8±3.1	4.2±2.9	4.3±3.0	*	4.5±3.1
Midland County	8.3±3.5	7.3±3.3	5.6±2.9	4.0±2.5	3.7±2.4	4.0±2.5	4.9±2.8	5.8±3.0	6.3±3.2	4.9±2.8

Table 16-B
White Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2004-2015

Health Department	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Mid-Michigan District	4.1±1.6	4.1±1.6	5.0±1.8	4.7±1.8	5.6±1.9	5.8±2.0	5.4±1.9	4.3±1.7	4.9±1.8	5.5±2.0
Monroe County	6.9±2.3	6.8±2.4	6.4±2.3	4.8±2.0	4.7±1.9	3.5±1.7	3.6±1.8	5.1±2.1	5.3±2.2	6.0±2.3
Muskegon County	4.5±1.8	5.0±1.9	5.7±2.0	6.4±2.2	6.5±2.2	6.3±2.2	7.1±2.4	6.9±2.3	6.0±2.2	4.8±1.9
Northwest Michigan Community Health	4.8±2.3	5.6±2.5	6.4±2.7	6.5±2.8	6.4±2.9	6.8±3.0	6.9±3.0	6.8±3.0	5.8±2.8	5.7±2.8
Oakland County	4.9±0.7	4.9±0.8	4.9±0.8	4.8±0.8	5.3±0.8	5.1±0.8	5.7±0.9	5.6±0.9	5.4±0.8	4.5±0.8
Ottawa County	5.0±1.4	5.0±1.4	5.3±1.4	4.6±1.4	4.9±1.4	5.3±1.5	5.6±1.5	6.2±1.6	5.2±1.4	5.5±1.5
Delta & Menominee	4.9±3.2	4.8±3.3	3.8±3.0	4.3±3.4	*	*	4.5±3.3	3.8±3.0	4.9±3.4	*
Saginaw County	5.0±1.8	5.7±2.0	6.6±2.2	6.4±2.2	4.7±1.9	2.9±1.5	2.9±1.5	4.2±1.8	6.2±2.2	6.1±2.2
Sanilac County	5.8±3.8	7.2±4.2	7.0±4.3	6.3±4.1	*	*	*	4.6±3.7	6.9±4.5	6.2±4.3
Shiawassee County	7.8±3.4	9.0±3.8	5.5±3.0	5.9±3.2	3.3±2.4	3.4±2.5	3.9±2.7	3.3±2.5	4.3±2.8	4.3±2.8
St Clair County	7.0±2.1	5.8±2.0	5.0±1.9	6.2±2.1	5.4±2.1	6.3±2.3	6.2±2.3	6.3±2.3	5.9±2.3	4.3±1.9
Tuscola County	7.8±3.9	6.3±3.6	3.9±2.9	4.0±3.0	6.1±3.8	6.9±4.1	7.0±4.1	5.0±3.5	4.9±3.4	4.8±3.3
Van Buren-Cass County	5.5±2.2	4.4±2.0	5.0±2.2	5.9±2.4	6.0±2.4	4.5±2.1	5.8±2.5	6.7±2.7	7.1±2.7	5.8±2.4
Washtenaw County	4.5±1.4	3.5±1.2	3.5±1.3	4.3±1.4	4.4±1.5	4.5±1.5	3.9±1.4	4.2±1.5	5.4±1.7	4.8±1.6
Wayne County	5.0±0.7	5.3±0.8	5.6±0.8	5.6±0.8	5.4±0.8	5.4±0.8	5.5±0.9	5.9±0.9	5.9±0.9	6.1±0.9
Western Upper Peninsula District	5.8±3.4	6.8±3.7	7.3±3.8	4.8±3.1	4.7±3.1	3.8±2.8	4.4±3.1	4.1±3.0	5.3±3.5	4.8±3.3

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 16-C
Black Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2004-2015

Health Department	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	16.6±1.0	16.4±1.0	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.6±0.9
Barry-Eaton District	*	*	*	25.2±18.4	28.0±19.1	26.9±18.4	*	22.7±16.6	22.4±16.4	29.2±17.9
Berrien County	11.9±5.5	16.9±6.5	14.9±6.0	16.7±6.4	16.0±6.2	19.5±6.9	18.6±6.9	17.3±6.7	16.5±6.7	13.3±6.1
Calhoun County	20.8±9.8	25.3±10.7	24.2±10.5	20.5±9.6	22.1±10.4	25.4±11.3	20.2±10.1	13.8±8.1	9.9±6.8	14.6±8.2
City Of Detroit	16.8±1.4	16.2±1.4	15.8±1.4	16.5±1.5	16.0±1.5	15.0±1.5	14.7±1.5	14.8±1.5	15.1±1.5	14.7±1.5
District #10	-	*	*	*	*	*	*	*	*	*
Genesee County	18.6±3.8	17.1±3.6	17.1±3.6	17.9±3.7	14.6±3.4	14.2±3.4	12.4±3.3	14.2±3.5	14.6±3.6	15.6±3.7
Ingham County	19.8±6.8	19.4±6.8	17.8±6.3	16.5±6.0	16.7±5.9	13.1±5.3	11.4±5.0	10.1±4.7	10.1±4.7	12.9±5.2
Jackson County	28.9±13.5	28.7±13.4	16.8±10.3	15.6±10.1	13.2±9.1	11.6±8.6	13.0±9.0	18.5±10.8	16.2±10.0	15.0±9.7
Kalamazoo County	16.0±6.1	16.8±6.2	16.8±6.2	19.2±6.5	19.5±6.5	18.3±6.4	17.6±6.3	15.5±5.9	14.2±5.7	15.3±5.9
Kent County	18.0±4.4	16.8±4.2	14.4±3.8	13.4±3.7	13.9±3.8	16.6±4.2	14.6±3.9	12.9±3.6	9.9±3.2	10.4±3.3
Macomb County	12.7±4.5	13.5±4.3	13.4±4.0	15.1±4.0	13.8±3.8	13.1±3.6	11.4±3.2	10.8±3.0	9.6±2.8	10.6±2.9
Mid-Michigan District	*	*	-	-	-	*	*	*	-	*
Muskegon County	13.4±6.0	9.8±5.1	9.0±4.9	8.2±4.6	13.2±5.9	14.9±6.3	16.4±6.8	12.8±6.0	14.9±6.7	19.7±7.7
Oakland County	16.7±3.3	17.4±3.4	15.4±3.1	14.7±3.0	11.9±2.7	10.8±2.5	10.6±2.4	11.1±2.5	13.7±2.8	14.6±2.9
Ottawa County	-	-	*	*	*	*	*	*	*	*
Saginaw County	18.4±5.7	19.0±5.8	15.3±5.2	19.2±5.8	17.4±5.6	19.2±6.1	13.4±5.1	14.3±5.2	12.2±5.0	15.9±5.7
St Clair County	*	*	*	*	*	*	*	*	*	*
Van Buren-Cass County	28.6±20.9	33.3±22.7	*	*	*	*	*	*	*	*
Washtenaw County	12.1±4.6	13.4±4.9	13.2±4.9	11.6±4.6	10.5±4.4	8.7±4.0	8.9±4.1	8.3±4.1	10.9±4.7	12.1±5.2
Wayne County	16.0±3.2	14.4±3.0	12.7±2.7	10.4±2.4	11.7±2.5	12.6±2.5	13.6±2.6	11.3±2.4	9.6±2.2	8.1±2.0

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 16-A
Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2004-2015

City	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	7.6±0.3	7.8±0.3	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.8±0.3
Ann Arbor	5.3±2.4	5.4±2.5	3.9±2.1	4.3±2.2	3.7±2.1	3.1±1.9	2.7±1.8	3.6±2.0	5.1±2.4	4.3±2.2
Battle Creek	9.2±3.7	11.3±4.0	12.4±4.3	11.0±4.1	10.6±4.1	11.3±4.3	11.0±4.3	9.9±3.8	7.9±3.4	8.6±3.5
Dearborn	4.8±1.9	5.6±2.0	7.7±2.3	6.9±2.3	7.0±2.3	6.3±2.2	7.4±2.4	7.5±2.4	6.4±2.2	6.2±2.2
Dearborn Heights	6.0±3.1	7.6±3.5	5.8±3.0	5.8±3.0	3.8±2.5	5.2±2.9	5.7±3.1	6.9±3.4	5.5±3.0	5.2±2.8
Detroit	15.0±1.2	14.7±1.2	14.3±1.2	14.8±1.3	14.4±1.3	13.6±1.3	13.6±1.3	13.6±1.3	13.9±1.3	13.5±1.3
Farmington Hills	7.1±3.2	7.7±3.4	7.4±3.3	7.0±3.2	6.4±3.1	6.4±3.1	6.9±3.3	6.7±3.2	6.7±3.2	6.1±3.0
Flint	15.3±3.0	12.2±2.7	11.6±2.7	12.6±2.9	12.6±3.0	12.1±3.0	11.6±3.0	11.0±2.8	10.9±2.8	10.7±2.8
Grand Rapids	8.6±1.7	8.3±1.7	10.0±1.9	8.3±1.8	8.6±1.8	8.5±1.8	9.4±1.9	8.0±1.7	6.6±1.6	5.9±1.5
Kalamazoo	8.3±3.2	9.5±3.4	9.2±3.3	10.5±3.5	11.0±3.6	10.3±3.6	9.7±3.5	6.0±2.5	6.0±2.5	7.5±2.8
Kentwood	8.1±3.6	6.8±3.3	4.2±2.6	3.9±2.6	3.4±2.4	3.4±2.4	4.5±2.7	5.2±2.9	5.1±2.9	6.0±3.1
Lansing	8.0±2.2	8.0±2.3	8.5±2.3	8.9±2.4	8.7±2.4	7.1±2.2	6.5±2.1	7.4±2.2	8.5±2.4	8.8±2.4
Livonia	4.4±2.5	3.0±2.1	5.4±2.8	5.2±2.8	4.8±2.7	2.4±1.9	2.8±2.1	5.5±2.9	5.4±2.8	4.8±2.6
Midland	8.0±4.7	7.4±4.6	5.9±4.1	5.1±3.8	*	*	*	6.3±4.1	7.7±4.5	7.6±4.5
Muskegon	7.3±3.8	6.7±3.6	7.5±3.9	7.5±3.9	7.2±3.9	7.8±4.1	10.5±4.8	12.6±5.0	12.6±5.0	13.4±5.2
Novi	*	*	5.3±3.3	5.3±3.3	6.6±3.7	3.3±2.7	3.8±2.8	4.3±3.0	6.1±3.5	5.0±3.1
Pontiac	13.1±3.6	12.9±3.7	12.5±3.7	14.4±4.1	13.8±4.0	11.9±3.8	8.5±3.2	9.4±3.3	11.4±3.6	14.0±4.0
Portage	6.6±3.7	7.7±4.0	4.4±3.1	9.2±4.5	8.6±4.3	7.5±4.1	4.1±3.0	5.8±3.6	5.8±3.6	5.2±3.4
Rochester Hills	5.6±2.9	4.1±2.5	6.2±3.1	5.0±2.8	4.8±2.8	3.5±2.5	4.0±2.6	5.2±3.0	4.4±2.7	4.0±2.6
Roseville	5.3±3.3	9.9±4.6	9.7±4.4	10.0±4.6	5.5±3.4	5.7±3.5	7.4±4.0	7.8±4.1	7.8±4.0	8.3±4.2
Royal Oak	3.2±2.4	3.7±2.6	4.0±2.6	4.4±2.7	5.6±3.0	6.4±3.2	8.2±3.7	7.1±3.5	7.6±3.6	4.0±2.6
Saginaw	11.1±3.7	13.0±4.0	11.5±3.8	14.6±4.3	13.2±4.2	13.7±4.4	9.6±3.7	12.1±4.2	12.7±4.3	15.0±4.7
Southfield	10.7±4.1	13.5±4.6	12.4±4.5	13.1±4.6	11.2±4.4	10.0±4.1	10.9±4.3	8.5±3.7	11.7±4.4	9.0±3.8
St Clair Shores	6.0±3.5	4.2±2.9	3.7±2.7	5.4±3.4	4.0±3.0	4.7±3.3	*	4.7±3.2	4.6±3.2	5.6±3.5
Sterling Heights	3.7±1.8	6.2±2.3	7.0±2.4	7.5±2.5	7.0±2.5	6.2±2.4	6.6±2.4	5.7±2.3	6.7±2.4	6.4±2.4
Taylor	10.8±4.0	10.2±3.9	8.5±3.5	6.3±3.1	6.8±3.2	9.9±3.9	11.8±4.2	10.8±4.0	7.4±3.3	5.8±2.9
Troy	4.4±2.6	4.9±2.8	5.8±3.0	5.5±3.0	6.2±3.2	4.4±2.7	3.8±2.5	3.3±2.3	3.8±2.4	4.1±2.5
Warren	7.6±2.4	7.2±2.3	6.4±2.2	8.1±2.5	7.9±2.5	9.1±2.7	7.6±2.4	8.7±2.6	6.6±2.2	5.4±2.0
Westland	6.4±2.7	6.3±2.7	7.5±2.9	7.4±2.9	8.2±3.1	9.6±3.4	10.4±3.5	9.3±3.3	8.3±3.1	7.4±2.9
Wyoming	8.9±3.0	10.9±3.2	12.1±3.4	11.0±3.3	8.4±3.0	6.7±2.6	4.6±2.2	4.1±2.1	4.4±2.1	6.0±2.5

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 17-B
White Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2004-2015

City	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	5.4±0.3	5.6±0.3	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3
Ann Arbor	3.0±2.2	3.2±2.3	*	3.7±2.5	3.2±2.4	3.7±2.6	*	3.2±2.3	4.5±2.8	4.0±2.6
Battle Creek	6.0±3.4	9.1±4.2	9.5±4.4	7.2±3.9	4.8±3.3	5.5±3.6	8.1±4.4	8.1±4.1	7.0±3.8	5.9±3.5
Dearborn	3.6±1.7	5.4±2.1	7.4±2.4	6.4±2.3	6.2±2.3	6.0±2.3	6.5±2.4	7.0±2.5	5.9±2.4	7.1±2.7
Dearborn Heights	6.3±3.4	7.2±3.6	5.2±3.1	4.3±2.8	3.0±2.4	3.1±2.5	3.7±2.7	5.3±3.2	6.0±3.5	6.9±3.7
Detroit	4.9±1.6	5.5±1.8	5.2±1.9	5.5±2.1	5.8±2.3	6.5±2.8	9.9±3.8	10.1±4.0	11.3±4.2	9.9±3.9
Farmington Hills	6.9±3.9	6.8±4.0	7.2±4.3	6.1±4.0	5.6±3.9	6.5±4.2	8.0±4.7	6.5±4.2	5.0±3.7	4.9±3.6
Flint	8.0±3.4	6.5±3.2	5.7±3.1	6.7±3.5	9.6±4.3	7.1±3.8	7.6±4.1	5.5±3.4	5.9±3.5	4.9±3.2
Grand Rapids	3.4±1.3	3.4±1.3	5.5±1.7	4.7±1.6	5.1±1.7	4.4±1.6	5.7±1.8	5.3±1.7	4.6±1.6	4.2±1.5
Kalamazoo	5.5±3.3	4.6±3.0	3.6±2.7	*	3.6±2.7	4.8±3.1	6.1±3.6	3.5±2.4	2.6±2.0	2.9±2.2
Kentwood	5.8±3.6	4.7±3.2	*	*	*	*	*	4.5±3.3	5.2±3.6	5.4±3.7
Lansing	3.4±1.8	3.8±1.9	4.5±2.1	5.2±2.3	4.6±2.2	5.2±2.4	5.3±2.5	7.4±2.9	7.9±3.1	7.5±3.1
Livonia	4.8±2.7	3.3±2.3	5.6±3.0	5.4±3.0	5.0±2.9	*	*	4.9±2.9	5.3±3.0	5.1±2.9
Midland	7.0±4.6	7.9±4.9	6.4±4.4	5.5±4.1	*	*	*	4.7±3.7	6.1±4.2	6.0±4.2
Muskegon	4.9±3.9	6.8±4.7	7.2±4.9	6.7±4.9	*	*	7.1±5.2	8.6±5.3	6.0±4.4	5.3±4.2
Novi	*	*	*	*	5.1±4.1	*	*	*	5.1±4.1	*
Pontiac	5.0±3.1	5.4±3.3	5.7±3.5	8.5±4.6	7.7±4.6	6.3±4.4	*	4.9±3.9	4.8±3.8	6.2±4.3
Portage	5.1±3.5	5.6±3.7	3.8±3.0	6.8±4.2	7.4±4.4	6.1±4.0	4.1±3.3	5.4±3.7	5.4±3.7	4.7±3.5
Rochester Hills	4.1±2.8	3.1±2.5	7.0±3.8	5.5±3.4	5.6±3.5	3.4±2.7	3.4±2.7	3.9±2.9	*	3.5±2.8
Roseville	4.4±3.2	8.5±4.6	9.8±4.9	10.3±5.2	6.3±4.1	5.3±3.9	5.6±4.1	4.7±3.8	6.3±4.3	5.5±4.1
Royal Oak	*	*	3.0±2.4	3.4±2.5	5.8±3.3	6.8±3.5	8.3±3.9	6.5±3.5	6.9±3.6	3.9±2.7
Saginaw	*	5.3±3.6	6.2±4.0	6.3±4.1	4.3±3.5	*	*	5.7±4.2	9.7±5.4	10.6±5.7
Southfield	*	*	*	*	11.6±9.2	11.9±9.5	12.0±9.5	*	*	-
St Clair Shores	5.2±3.4	4.0±2.9	4.1±3.0	6.0±3.7	4.5±3.4	5.3±3.6	*	*	*	4.9±3.4
Sterling Heights	4.0±2.0	5.7±2.4	6.1±2.5	6.9±2.7	7.7±2.9	6.9±2.8	7.6±3.0	6.0±2.6	6.5±2.7	5.4±2.4
Taylor	9.7±4.3	7.9±4.0	7.0±3.8	5.6±3.5	5.3±3.5	6.7±4.0	8.2±4.4	8.0±4.3	6.8±4.0	6.1±3.8
Troy	3.5±2.8	4.9±3.4	5.1±3.5	5.3±3.6	6.1±4.0	5.4±3.7	5.9±3.9	4.6±3.4	4.7±3.4	3.9±3.1
Warren	6.5±2.5	7.1±2.6	5.9±2.5	6.8±2.7	6.3±2.7	6.9±2.9	5.9±2.7	8.0±3.2	6.7±2.9	5.7±2.7
Westland	3.4±2.2	4.4±2.6	4.9±2.8	5.9±3.1	5.7±3.1	7.8±3.7	8.6±3.9	8.5±3.9	9.0±4.0	8.0±3.8
Wyoming	7.4±2.9	9.1±3.2	9.3±3.2	8.6±3.2	6.7±2.8	5.3±2.5	3.8±2.1	3.2±2.0	4.2±2.3	5.7±2.7

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 17-C
Black Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2004-2015

City	2004-06	2005-07	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15
Michigan	16.6±1.0	16.4±1.0	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.6±0.9
Battle Creek	18.2±11.2	19.1±11.2	20.9±11.7	21.0±11.7	23.0±12.9	21.5±12.6	11.9±9.5	12.1±8.9	11.6±8.5	15.9±9.8
Detroit	16.8±1.4	16.2±1.4	15.8±1.4	16.5±1.5	15.9±1.5	15.0±1.5	14.7±1.5	14.8±1.5	15.0±1.5	14.7±1.5
Flint	20.1±4.6	15.8±4.1	15.5±4.1	16.6±4.2	14.7±4.1	15.3±4.3	14.1±4.2	14.7±4.2	14.4±4.2	14.7±4.3
Grand Rapids	17.0±5.1	17.7±5.1	17.6±5.1	16.8±5.0	16.0±5.0	18.8±5.4	16.7±5.1	14.9±4.8	11.5±4.2	12.0±4.4
Kalamazoo	14.0±7.0	17.4±7.7	18.5±8.0	22.4±8.7	21.6±8.5	17.7±7.9	12.7±6.9	9.7±5.7	11.5±6.2	16.7±7.4
Lansing	20.9±7.3	20.0±7.1	19.0±6.7	17.9±6.4	18.3±6.5	13.5±5.7	11.0±5.2	8.9±4.7	10.7±5.1	13.5±5.6
Muskegon	11.8±8.1	*	*	*	12.0±8.3	13.1±8.5	17.4±10.2	18.1±10.1	20.1±10.8	21.7±11.2
Pontiac	22.6±7.1	21.6±7.1	19.6±6.7	19.9±6.8	19.1±6.7	16.5±6.2	12.5±5.3	12.6±5.2	16.4±5.9	19.4±6.5
Saginaw	18.6±6.7	20.2±7.0	16.0±6.2	20.1±7.0	18.4±6.9	20.5±7.5	14.1±6.3	15.8±6.5	14.2±6.3	17.3±7.2
Southfield	12.9±5.3	15.4±5.9	14.8±5.8	15.0±5.8	11.6±5.2	10.0±4.7	11.2±5.0	10.3±4.7	14.9±5.8	12.3±5.2
Taylor	17.0±11.0	19.9±11.6	14.9±9.7	9.3±7.4	11.9±8.2	18.8±9.8	21.6±10.2	18.7±9.4	10.2±7.0	*
Warren	13.1±9.0	9.8±7.3	11.8±7.2	14.8±7.7	14.6±7.3	14.1±6.9	11.9±6.0	9.4±5.1	7.8±4.6	5.7±3.9
Westland	19.5±12.0	10.6±8.5	12.3±8.5	11.0±7.6	14.0±8.2	12.1±7.5	14.7±8.3	11.7±7.2	9.1±6.3	6.6±5.3

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2004-2013 Geocoded Michigan Death Certificate Registries; 2014-2015 Michigan Death Certificate Registries. 2004-2015 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 18-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	700	660	697	608	600	549	501	540	544	513	498
Alcona	1	-	-	1	1	-	-	1	-	1	-
Alger	-	-	-	3	-	-	-	-	-	-	-
Allegan	10	4	13	4	4	6	5	3	3	4	9
Alpena	-	3	-	1	-	-	1	-	1	1	1
Antrim	-	1	-	2	-	4	2	-	3	-	-
Arenac	-	1	-	-	2	-	1	1	4	2	-
Baraga	2	-	-	1	-	-	-	-	1	1	1
Barry	1	6	3	2	3	4	2	2	3	1	-
Bay	8	8	5	3	7	9	6	4	-	5	4
Benzie	1	2	2	-	-	-	-	-	2	-	1
Berrien	10	12	13	10	10	8	8	11	8	9	6
Branch	2	2	2	1	2	2	3	3	2	3	2
Calhoun	11	11	18	13	13	10	9	7	8	5	8
Cass	6	1	2	2	5	1	2	2	-	5	-
Charlevoix	1	-	2	1	2	-	-	-	-	-	2
Cheboygan	-	-	1	-	-	-	-	-	-	-	-
Chippewa	4	2	-	1	-	-	3	2	1	1	-
Clare	-	2	3	-	1	-	1	2	1	1	-
Clinton	2	2	6	1	1	3	2	2	1	3	5
Crawford	-	2	4	1	1	-	-	-	1	-	-
Delta	-	-	2	-	2	2	-	4	1	3	-
Dickinson	-	1	-	2	2	1	-	-	4	1	-
Eaton	3	4	4	3	3	7	5	2	6	3	7
Emmet	1	-	1	2	-	3	-	1	2	1	-
Genesee	33	44	40	27	34	19	22	19	25	23	29

**Table 18-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015**

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Gladwin	4	–	1	1	1	1	1	–	1	1	2
Gogebic	–	1	1	–	–	2	–	–	–	1	–
Grand Traverse	2	4	5	7	3	3	–	3	–	1	7
Gratiot	–	4	4	2	3	3	3	–	3	2	–
Hillsdale	–	6	4	3	1	1	2	1	3	–	3
Houghton	1	2	2	3	1	3	1	2	2	–	–
Huron	2	2	–	–	3	1	1	–	–	–	–
Ingham	19	13	17	22	14	12	16	16	20	15	11
Ionia	2	3	2	–	1	2	1	–	2	2	2
Iosco	1	1	–	–	1	1	1	–	–	2	1
Iron	–	–	1	–	–	1	–	–	–	–	–
Isabella	7	3	5	1	1	–	6	2	–	2	4
Jackson	17	11	12	11	7	9	8	3	10	5	7
Kalamazoo	16	14	22	13	20	17	8	17	11	11	11
Kalkaska	3	–	–	–	1	1	–	3	1	3	–
Kent	45	59	47	48	42	32	40	38	28	37	38
Keweenaw	–	1	–	–	–	–	–	–	–	–	–
Lake	2	–	–	–	1	–	–	1	–	–	1
Lapeer	–	5	4	3	6	4	–	1	4	3	1
Leelanau	1	–	–	–	–	3	–	–	–	1	–
Lenawee	6	8	5	9	8	11	3	6	4	5	4
Livingston	8	5	11	6	9	6	9	7	3	9	5
Luce	1	1	–	1	–	–	–	–	–	1	–
Mackinac	1	1	–	1	1	–	–	–	–	–	2
Macomb	41	30	48	40	52	39	43	40	52	34	43

Table 18-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Manistee	1	1	-	1	1	1	-	-	-	1	-
Marquette	2	2	2	2	2	1	3	-	2	-	1
Mason	1	3	-	1	1	2	1	-	-	1	2
Mecosta	1	2	1	4	-	1	4	4	4	2	3
Menominee	2	-	1	1	1	-	1	-	-	1	1
Midland	7	7	3	1	3	2	2	3	4	3	-
Missaukee	1	-	-	-	1	-	-	1	-	2	-
Monroe	7	11	6	5	3	4	2	6	10	2	8
Montcalm	-	1	2	1	2	5	3	3	4	1	2
Montmorency	-	1	-	-	-	1	-	-	-	-	-
Muskegon	10	9	13	9	9	16	11	11	14	9	7
Newaygo	5	1	6	3	1	2	6	2	2	3	-
Oakland	73	72	71	62	60	63	45	61	69	66	53
Oceana	3	1	1	2	2	3	-	2	1	3	1
Ogemaw	-	-	1	-	-	1	1	1	-	-	-
Ontonagon	-	1	1	-	-	-	-	-	1	1	-
Osceola	3	2	1	-	1	1	-	-	1	-	2
Oscoda	-	-	-	1	1	-	-	1	-	1	-
Otsego	-	-	3	1	-	2	3	1	-	1	1
Ottawa	11	18	16	19	10	17	17	14	20	10	12
Presque Isle	-	1	2	-	-	1	1	3	-	1	2
Roscommon	1	-	3	1	1	-	1	-	-	1	1
Saginaw	18	9	16	10	15	10	8	8	13	11	12
St. Clair	10	9	10	4	6	4	6	10	2	3	6
St. Joseph	4	7	6	8	9	2	3	1	5	3	1

Table 18-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Sanilac	1	4	4	1	2	-	-	1	1	3	3
Schoolcraft	1	-	1	-	-	-	-	-	-	-	-
Shiawassee	4	-	8	-	2	2	1	2	2	1	1
Tuscola	3	3	1	-	2	4	1	4	1	-	3
Van Buren	9	-	5	6	4	2	3	4	4	3	3
Washtenaw	19	15	17	14	18	11	10	14	12	23	10
Wayne	225	198	180	197	172	158	151	172	147	147	144
Detroit City	161	121	120	123	114	98	80	104	87	80	81
Wayne County Excluding Detroit City	64	77	60	74	58	60	71	68	60	67	63
Wexford	2	-	4	2	2	2	2	5	3	2	1
Unknown	1	-	-	-	-	-	-	-	1	-	1

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Neonatal Death: Deaths occurring to individuals less than 28 days of age.
 -: A dash indicates quantity zero.

Source: 2005-2015 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 18-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	313	280	300	286	281	268	248	243	255	260	267
Alcona	-	1	-	-	-	-	1	-	-	-	-
Alger	-	-	-	-	1	-	-	-	-	-	-
Allegan	3	4	5	2	2	-	2	1	6	5	1
Alpena	-	1	1	-	3	1	1	-	-	1	-
Antrim	-	1	2	-	1	-	-	-	-	-	-
Arenac	-	-	-	-	-	1	1	-	-	-	-
Baraga	-	-	-	-	-	-	-	-	-	-	-
Barry	1	1	2	3	2	4	-	1	1	1	-
Bay	2	4	3	5	1	3	3	1	2	5	2
Benzie	2	-	-	-	-	1	-	-	-	-	-
Berrien	10	5	8	6	7	6	10	3	6	6	3
Branch	4	-	-	1	-	2	1	1	1	1	1
Calhoun	7	6	10	1	7	8	6	3	3	4	6
Cass	2	2	1	2	1	1	1	2	3	-	3
Charlevoix	-	1	-	-	-	-	1	-	-	1	1
Cheboygan	-	1	1	1	2	-	-	-	-	-	1
Chippewa	1	2	1	-	-	2	1	-	1	1	-
Clare	-	1	-	1	1	1	-	-	-	-	1
Clinton	1	1	-	2	1	2	-	-	1	2	3
Crawford	-	-	1	-	-	-	1	-	-	-	2
Delta	1	1	2	-	1	-	1	-	-	-	1
Dickinson	2	3	-	-	-	1	-	-	-	-	-
Eaton	1	2	2	4	3	1	1	3	4	3	2
Emmet	2	-	-	-	1	-	-	1	-	-	-
Genesee	20	17	17	19	17	11	18	19	15	18	14

**Table 18-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015**

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Gladwin	-	-	-	-	1	1	-	-	1	2	2
Gogebic	-	1	-	-	-	-	-	-	-	-	-
Grand Traverse	3	2	2	1	2	-	1	1	-	5	2
Gratiot	1	1	-	1	-	2	-	-	-	2	1
Hillsdale	-	1	1	1	-	1	-	1	-	-	2
Houghton	-	-	1	1	-	-	-	-	1	1	-
Huron	-	1	-	-	-	-	1	3	-	-	-
Ingham	11	5	9	3	6	2	5	6	8	6	10
Ionia	1	-	4	1	-	3	2	1	3	1	1
Iosco	-	-	1	1	2	2	1	1	1	-	-
Iron	-	-	-	-	1	1	1	-	-	1	-
Isabella	2	3	2	-	3	4	-	2	2	-	1
Jackson	8	6	3	3	5	2	3	4	5	3	3
Kalamazoo	6	3	7	9	9	13	3	2	15	5	9
Kalkaska	-	-	-	-	-	1	-	1	1	-	-
Kent	21	26	18	21	14	22	19	12	11	15	18
Keweenaw	-	-	-	-	-	-	-	-	-	-	-
Lake	1	1	-	-	1	1	-	-	-	-	-
Lapeer	3	1	1	1	5	3	-	1	-	2	3
Leelanau	-	-	1	-	-	-	1	-	1	2	1
Lenawee	-	2	4	2	1	4	3	2	5	4	2
Livingston	2	2	3	4	3	5	3	1	4	4	2
Luce	-	1	-	1	-	-	-	-	-	1	1
Mackinac	-	-	-	1	-	-	-	-	-	-	-
Macomb	20	21	21	22	26	9	10	12	15	14	17

**Table 18-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015**

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Manistee	3	-	-	-	1	1	1	-	-	1	-
Marquette	1	2	-	-	2	1	1	2	-	1	4
Mason	-	1	1	1	-	1	-	-	-	1	-
Mecosta	1	-	1	5	-	2	2	-	1	1	-
Menominee	-	-	-	1	-	-	-	1	-	-	-
Midland	1	1	-	2	2	-	2	3	3	2	3
Missaukee	1	-	-	-	-	-	-	-	-	-	-
Monroe	4	4	4	6	6	5	1	3	5	1	1
Montcalm	1	3	-	2	2	1	1	2	1	1	1
Montmorency	-	-	-	-	-	-	-	2	-	-	-
Muskegon	3	4	6	7	6	7	4	7	5	6	11
Newaygo	1	3	3	-	2	1	1	-	1	4	3
Oakland	30	19	21	26	17	21	25	32	21	17	18
Oceana	3	1	1	1	-	1	-	3	-	3	1
Ogemaw	1	-	1	1	1	1	-	1	-	1	-
Ontonagon	-	-	-	-	-	-	-	-	-	-	-
Osceola	1	-	1	-	-	1	2	-	-	-	-
Oscoda	-	-	-	1	-	-	-	1	-	-	-
Otsego	-	2	2	-	1	-	1	2	3	2	-
Ottawa	5	4	1	4	5	3	4	4	6	2	9
Presque Isle	-	-	-	-	1	1	-	-	-	-	-
Roscommon	-	-	1	-	-	-	-	-	-	-	-
Saginaw	10	10	10	15	11	2	9	5	6	10	7
St. Clair	4	4	2	4	9	2	5	3	4	6	3
St. Joseph	2	2	4	2	1	5	1	6	2	2	-

Table 18-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2005-2015

County	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Sanilac	1	-	2	-	1	1	-	3	1	-	1
Schoolcraft	-	1	-	-	-	-	-	-	-	-	-
Shiawassee	6	2	2	1	-	2	-	1	1	2	3
Tuscola	3	1	1	1	2	1	1	-	1	2	1
Van Buren	5	1	1	4	1	2	1	4	2	-	3
Washtenaw	6	8	6	3	4	9	4	6	6	5	4
Wayne	78	75	94	78	76	75	79	66	70	71	78
Detroit City	49	54	59	52	52	48	50	47	47	50	58
Wayne County Excluding Detroit City	29	21	35	26	24	27	29	19	23	21	20
Wexford	4	1	1	1	-	1	1	1	-	2	-
Unknown	-	-	-	-	-	-	-	-	-	1	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Postneonatal Death: Deaths occurring to individuals 28–364 days of age.
 -: A dash indicates quantity zero.

Source: 2005-2015 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 19-A
Number of Neonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2005-2015**

Health Department	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	700	660	697	608	600	549	501	540	544	513	498
Allegan County	10	4	13	4	4	6	5	3	3	4	9
Barry-Eaton District	4	10	7	5	6	11	7	4	9	4	7
Berrien County	10	12	13	10	10	8	8	11	8	9	6
Branch-Hillsdale-St. Joseph	6	15	12	12	12	5	8	5	10	6	6
Calhoun County	11	11	18	13	13	10	9	7	8	5	8
Central Michigan District	15	8	13	3	7	2	10	5	7	7	9
Chippewa County	4	2	-	1	-	-	3	2	1	1	-
City Of Detroit	161	121	120	123	114	98	80	104	87	80	81
Delta & Menominee	2	-	3	1	3	2	1	4	1	4	1
Dickinson-Iron District	-	1	1	2	2	2	-	-	4	1	-
District #10	19	10	16	14	11	12	13	18	12	17	8
District #2	2	1	1	2	3	2	2	3	-	4	1
District #4	-	5	3	1	-	2	2	3	1	2	3
Genesee County	33	44	40	27	34	19	22	19	25	23	29
Grand Traverse County	2	4	5	7	3	3	-	3	-	1	7
Ingham County	19	13	17	22	14	12	16	16	20	15	11
Ionia County	2	3	2	-	1	2	1	-	2	2	2
Jackson County	17	11	12	11	7	9	8	3	10	5	7
Kalamazoo County	16	14	22	13	20	17	8	17	11	11	11
Kent County	45	59	47	48	42	32	40	38	28	37	38
Lapeer County	-	5	4	3	6	4	-	1	4	3	1
Lenawee County	6	8	5	9	8	11	3	6	4	5	4
Livingston County	8	5	11	6	9	6	9	7	3	9	5
Luce-Mackinac-Alger-Schoolcraft	3	2	1	5	1	-	-	-	-	1	2
Macomb County	41	30	48	40	52	39	43	40	52	34	43

Table 19-A
Number of Neonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2005-2015

Health Department	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Marquette County	2	2	2	2	2	1	3	-	2	-	1
Mid-Michigan District	2	7	12	4	6	11	8	5	8	6	7
Midland County	7	7	3	1	3	2	2	3	4	3	-
Monroe County	7	11	6	5	3	4	2	6	10	2	8
Muskegon County	10	9	13	9	9	16	11	11	14	9	7
Oakland County	73	72	71	62	60	63	45	61	69	66	53
Ottawa County	11	18	16	19	10	17	17	14	20	10	12
Bay County	8	8	5	3	7	9	6	4	-	5	4
Saginaw County	18	9	16	10	15	10	8	8	13	11	12
Shiawassee County	4	-	8	-	2	2	1	2	2	1	1
St Clair County	10	9	10	4	6	4	6	10	2	3	6
Van Buren-Cass County	15	1	7	8	9	3	5	6	4	8	3
Washtenaw County	19	15	17	14	18	11	10	14	12	23	10
Wayne County	64	77	60	74	58	60	71	68	60	67	63
Benzie-Leelanau District	2	2	2	-	-	3	-	-	2	1	1
Northwest Michigan Community Health	2	1	6	6	2	9	5	2	5	2	3
Western Upper Peninsula District	3	5	4	4	1	5	1	2	4	3	1
Tuscola County	3	3	1	-	2	4	1	4	1	-	3
Huron County	2	2	-	-	3	1	1	-	-	-	-
Sanilac County	1	4	4	1	2	-	-	1	1	3	3
Unknown	1	-	-	-	-	-	-	-	1	-	1

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Neonatal Death: Deaths occurring to individuals less than 28 days of age.
 -: A dash indicates quantity zero.

Source: 2005-2015 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 19-B
Number of Postneonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2005-2015

Health Department	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Michigan	313	280	300	286	281	268	248	243	255	260	267
Allegan County	3	4	5	2	2	-	2	1	6	5	1
Barry-Eaton District	2	3	4	7	5	5	1	4	5	4	2
Berrien County	10	5	8	6	7	6	10	3	6	6	3
Branch-Hillsdale-St. Joseph	6	3	5	4	1	8	2	8	3	3	3
Calhoun County	7	6	10	1	7	8	6	3	3	4	6
Central Michigan District	3	4	4	1	5	8	3	2	3	2	4
Chippewa County	1	2	1	-	-	2	1	-	1	1	-
City Of Detroit	49	54	59	52	52	48	50	47	47	50	58
Delta & Menominee	1	1	2	1	1	-	1	1	-	-	1
Dickinson-Iron District	2	3	-	-	1	2	1	-	-	1	-
District #10	14	7	8	8	4	9	6	5	3	12	6
District #2	1	1	2	3	3	3	2	3	1	1	-
District #4	-	2	2	1	6	2	1	2	-	1	1
Genesee County	20	17	17	19	17	11	18	19	15	18	14
Grand Traverse County	3	2	2	1	2	-	1	1	-	5	2
Ingham County	11	5	9	3	6	2	5	6	8	6	10
Ionia County	1	-	4	1	-	3	2	1	3	1	1
Jackson County	8	6	3	3	5	2	3	4	5	3	3
Kalamazoo County	6	3	7	9	9	13	3	2	15	5	9
Kent County	21	26	18	21	14	22	19	12	11	15	18
Lapeer County	3	1	1	1	5	3	-	1	-	2	3
Lenawee County	-	2	4	2	1	4	3	2	5	4	2
Livingston County	2	2	3	4	3	5	3	1	4	4	2
Luce-Mackinac-Alger-Schoolcraft	-	2	-	2	1	-	-	-	-	1	1
Macomb County	20	21	21	22	26	9	10	12	15	14	17

Table 19-B
Number of Postneonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2005-2015

Health Department	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Marquette County	1	2	-	-	2	1	1	2	-	1	4
Mid-Michigan District	3	5	-	5	3	5	1	2	2	5	5
Midland County	1	1	-	2	2	-	2	3	3	2	3
Monroe County	4	4	4	6	6	5	1	3	5	1	1
Muskegon County	3	4	6	7	6	7	4	7	5	6	11
Oakland County	30	19	21	26	17	21	25	32	21	17	18
Ottawa County	5	4	1	4	5	3	4	4	6	2	9
Bay County	2	4	3	5	1	3	3	1	2	5	2
Saginaw County	10	10	10	15	11	2	9	5	6	10	7
Shiawassee County	6	2	2	1	-	2	-	1	1	2	3
St Clair County	4	4	2	4	9	2	5	3	4	6	3
Van Buren-Cass County	7	3	2	6	2	3	2	6	5	-	6
Washtenaw County	6	8	6	3	4	9	4	6	6	5	4
Wayne County	29	21	35	26	24	27	29	19	23	21	20
Benzie-Leelanau District	2	-	1	-	-	1	1	-	1	2	1
Northwest Michigan Community Health	2	4	4	-	3	-	2	3	3	3	1
Western Upper Peninsula District	-	1	1	1	-	-	-	-	1	1	-
Tuscola County	3	1	1	1	2	1	1	-	1	2	1
Huron County	-	1	-	-	-	-	1	3	-	-	-
Sanilac County	1	-	2	-	1	1	-	3	1	-	1
Unknown	-	-	-	-	-	-	-	-	-	1	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Postneonatal Death: Deaths occurring to individuals 28-364 days of age.
 -: A dash indicates quantity zero.

Source: 2005-2015 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services